

Sets and Functions

A **set** is a group of numbers which share common properties. Some common sets are:

Natural Numbers

$$N = \{1, 2, 3, 4, 5, \dots\}$$

Whole Numbers

$$W = \{0, 1, 2, 3, 4, 5, \dots\}$$

Integers

$$Z = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$$

Rational Numbers

Q = all integers **and** fractions of them (e.g. $\frac{3}{4}$, $-\frac{5}{8}$, etc)


Real Numbers

R = all rational **and** irrational numbers (e.g. $\sqrt{2}$, π , etc.)


Sets are written inside curly brackets. The set with no members “{ }” is called the **empty set**.

\in means “is a member of”, e.g. $5 \in \{3, 4, 5, 6, 7\}$ \notin means “is not a member of”, e.g. $5 \notin \{6, 7, 8\}$

A **function** is a rule which links an element in Set A to **one and only one** element in Set B.




This shows a function



This does **not** show a function


Example 1: Each function below is defined on the set of real numbers. State the range of each.

a) $f(x) = \sin x$




$-1 \leq f(x) \leq 1$

b) $g(x) = x^2$



$g(x) \geq 0$

c) $h(x) = 1 - x^2$



$h(x) \leq 1$

When choosing the domain, two cases
MUST be avoided:

a) Denominators can't be zero

b) Can't find the square root of a negative value

e.g. For $f(x) = \frac{1}{x+5}$, $x \neq -5$, i.e. $\{x \in \mathbb{R} : x \neq -5\}$

e.g. For $g(x) = \sqrt{x-3}$, $x \geq 3$, i.e. $\{x \in \mathbb{R} : x \geq 3\}$

Example 2: For each function, state a suitable domain.

a) $g(x) = \sqrt{3x-2}$

$$\begin{aligned} 3x-2 &\geq 0 \\ 3x &\geq 2 \\ x &\geq \frac{2}{3} \\ &= \end{aligned}$$

b) $p(\theta) = \frac{2}{5-\theta}$

$$\begin{aligned} 5-\theta &\neq 0 \\ -\theta &\neq -5 \\ \theta &\neq 5 \\ &= \end{aligned}$$

c) $f(y) = \frac{y^2}{\sqrt{y-1}}$

$$\begin{aligned} y-1 &> 0 \\ y &> 1 \\ &= \end{aligned}$$

Composite Functions

In the linear function $y = 3x - 5$, we get y by doing two acts: (i) multiply x by 3; (ii) then subtract 5. This is called a **composite function**, where we “do” a function to the range of another function.

e.g. If $h(x)$ is the composite function obtained by performing $f(x)$ on $g(x)$, then we say

$$h(x) = f(g(x)) \text{ (“f of g of x”)}$$

Example 3: $f(x) = 5x + 1$ and $g(x) = 3x^2 + 2x$.

a) Find $f(g(-1))$

$$\begin{aligned} g(-1) &= 3(-1)^2 + 2(-1) \\ &= 3 + (-2) \\ &= 1 \\ f(1) &= 5(1) + 1 \\ &= \underline{\underline{6}}. \end{aligned}$$

c) Find $f(f(x))$

$$\begin{aligned} f(5x+1) &= 5(5x+1) + 1 \\ &= 25x + 5 + 1 \\ &= \underline{\underline{25x + 6}}. \end{aligned}$$

b) Find $f(g(x))$

$$\begin{aligned} &= f(3x^2 + 2x) \\ &= 5(3x^2 + 2x) + 1 \\ &= \underline{\underline{15x^2 + 10x + 1}} \end{aligned}$$

d) Find $g(f(x))$

$$\begin{aligned} &= g(5x+1) \\ &= 3(5x+1)^2 + 2(5x+1) \\ &= 3(25x^2 + 10x + 1) + 10x + 2 \\ &= 75x^2 + 30x + 3 + 10x + 2 \\ &= \underline{\underline{75x^2 + 40x + 5}} \end{aligned}$$

NOTE: Usually, $f(g(x))$ and $g(f(x))$ are NOT the same!

Example 4: $f(x) = 2x + 1$, $g(x) = x^2 + 6$

a) Find formulae for:

$$\begin{aligned} \text{(i) } f(g(x)) &= 2(x^2 + 6) + 1 \\ &= 2x^2 + 12 + 1 \\ &= 2x^2 + 13 \end{aligned}$$

(ii) $g(f(x))$

$$\begin{aligned} &(2x + 1)^2 + 6 \\ &= 4x^2 + 4x + 7 \end{aligned}$$

b) Solve the equation $f(g(x)) = g(f(x))$

$$\begin{aligned} 2x^2 + 13 &= 4x^2 + 4x + 7 \\ 2x^2 + 4x - 6 &= 0 \\ 2(x^2 + 2x - 3) &= 0, \quad x \neq 3 \\ 2(x + 3)(x - 1) &= 0 \\ \underline{x = -3, x = 1} \end{aligned}$$

Example 5: $f(x) = \frac{3}{x+1}$, $x \neq -1$. Find an expression for $f(f(x))$, as a fraction in its simplest form.

$$\begin{aligned} f(f(x)) &= f\left(\frac{3}{x+1}\right) \\ &= \frac{3}{\frac{3}{x+1} + 1} \\ &= \frac{3}{\frac{4+x}{x+1}} \\ &= 3 \times \frac{x+1}{4+x} \\ &= \frac{3x+3}{4+x} \end{aligned}$$

$$\begin{aligned} &\frac{3}{x+1} + \frac{x+1}{x+1} \\ &= \frac{3+x+1}{x+1} \\ &= \frac{4+x}{x+1} \end{aligned}$$

Past Paper Example: Functions f and g are defined on a set of real numbers by

$$f(x) = x^2 + 3$$

$$g(x) = x + 4$$


a) Find expressions for:

(i) $f(g(x))$


(ii) $g(f(x))$

b) Show that $f(g(x)) + g(f(x)) = 0$ has no real roots

Example 1: Sketch and annotate the graph of $y = x^2 - 2x - 15$



Example 2: Sketch and annotate the graph of $y = x^2 - 4x + 4$




Example 3: in the spaces provided, make a **basic** sketch of the graph(s) of the function(s) stated.

a) $y = 2x + 1$

b) $3x + 4y - 12 = 0$

c) $y = -1$ and $x = 5$




d) $y = x^2$ and $y = 4$


e) $y = x^2 - 4$

f) $y = (x - 2)^2$ and $y = 2x - x^2$


Example 4: Sketch and annotate the graph of
 $y = x^2 - 2x - 8$



Example 5: Sketch and annotate the graph of
 $y = (x + 3)^2 + 1$



Example 6: Sketch the graphs of $y = \sin x^\circ$, $y = \cos x^\circ$ and $y = \tan x^\circ$ below.



For trig graphs, how soon the graph repeats itself horizontally is known as the **period**, and half of the vertical height is known as the **amplitude**.

Function	Period	Amplitude
$y = \sin x^\circ$	360°	1
$y = \cos x^\circ$	360°	1
$y = \tan x^\circ$	180°	∞

For the graphs of:




$y = a \sin bx^\circ + c$
and
 $y = a \cos bx^\circ + c$:

a = amplitude
 b = waves in 360°
 c = vertical shift



$y = a \tan bx^\circ + c$:

b = "waves" in 180°
 c = vertical shift

Example 7: Sketch the graphs of:




c) $y = -3\sin 3x^\circ - 2$



Compound Angles

A compound angle is one containing two parts, e.g. $(x - 60)^\circ$. The graphs of compound angles can be thought of as the trig version of $y = f(x - a)$, i.e. shifted left or right by a units.




Example 8: On the axes opposite, sketch:

a) $y = \sin x^\circ$

b) $y = \sin (x - 45)^\circ$

$$y = f(x) + 2 \quad y = f(x) + a \quad y = f(x) - 2$$



$y = f(x) + a$ is obtained by sliding $y = f(x)$:
 Vertically upwards if $a > 0$
 Vertically downwards if $a < 0$

Lesson Starter - 5B1 - Wed 13/6/18

1) Let $f(x) = 3x^2 + 9$ & $g(x) = 2x - 1$ find:

a) $f(g(x))$	b) $g(f(x))$	c) $g(g(x))$
$= 3(2x-1)^2 + 9$	$= 2(3x^2+9) - 1$	$= 2(2x-1) - 1$
$= 3(4x^2 - 4x + 1) + 9$	$= 6x^2 + 18 - 1$	$= 4x - 2 - 1$
$= 12x^2 - 12x + 3 + 9$	$= \underline{\underline{6x^2 + 17}}$	$= \underline{\underline{4x - 3}}$
$= \underline{\underline{12x^2 - 12x + 12}}$		

2) Let $k(x) = \frac{x}{1-x}$ find $k^{-1}(x)$

$$y = \frac{x}{1-x}$$

$$y(1-x) = x$$


$$y - yx = x$$


$$x + yx = y$$


$$x(1+y) = y$$


$$x = \frac{y}{1+y}$$


$$k^{-1}(x) = \underline{\underline{\frac{x}{1+x}}}$$














Example 9: Part of the graph of $y = f(x)$ is shown.

On separate diagrams, sketch:

a) $y = f(-x) + 2$

b) $y = -\frac{1}{2} f(x+1)$



Past Paper Example: The diagram shows a sketch of the function $y = f(x)$.


To the diagram, add the graphs of:

a) $y = f(2x)$

b) $y = 1 - f(2x)$. $y = -f(2x) + 1$

($-4, 8$) \rightarrow ($-2, 8$)
($0, 0$) \rightarrow ($0, 0$)
($2, 8$) \rightarrow ($1, 8$)

($-4, 8$) \rightarrow ($-4, -8$) \rightarrow ($-2, -8$) \rightarrow ($-2, 2$)
($0, 0$) \rightarrow ($0, 0$) \rightarrow ($0, 2$) \rightarrow ($0, 1$)
($2, 8$) \rightarrow ($2, -8$) \rightarrow ($1, -8$) \rightarrow ($1, 2$)



Example 1: A sequence is defined by the recurrence relation $U_{n+1} = 3U_n + 2$, $U_0 = 4$.

Find the value of U_4 .

Example 2: A sequence is defined by the recurrence relation $U_n = 4U_{n-1} - 3$, where $U_0 = a$.

Find an expression for U_2 in terms of a .

Finding a Formula

Recurrence relations can be used to describe situations seen in real life where a quantity changes by the same percentage at regular intervals. The first thing to do in most cases is find a formula to describe the situation.

Example: Jennifer puts £5000 into a high-interest savings account which pays 7.5% p/a. Find a recurrence relation for the amount of money in the savings account.

Solution: Starting amount = £5000
 After 1 year: amount = starting amount + 7.5% (i.e. 107.5% of starting amount)
 = 1.075 x starting amount

Recurrence relation is: $U_{n+1} = 1.075U_n$ ($U_0 = 5000$)

Example 3: Find a recurrence relation to describe:

- | | |
|---|---|
| <p>a) The amount left to pay on a loan of £10000, with interest charged at 1.5% per month and fixed monthly payments of £250.</p> | <p>b) The amount of water in a swimming pool of volume 750,000 litres if 0.05% per day is lost to evaporation, but 350 litres extra is added daily.</p> |
|---|---|

Section 2 - Recurrence Relations.Additional Example 2.1.1

The value of a car depreciates by 5% per annum. Its value at the beginning of 2009 was £24,000.

- a) Find a recurrence relation for the value of the car.

$$\underline{U_{n+1} = 0.95U_n} \quad U_0 = \underline{24000}$$

- b) Calculate the expected value of the car at the beginning of 2013

$$U_1 = 0.95 \times 24000$$

$$U_1 = 22800$$

$$U_2 = 0.95 \times 22800$$

$$U_2 = 21660$$

$$U_3 = 0.95 \times 21660$$

$$U_3 = 20577$$

$$U_4 = 0.95 \times 20577$$

$$U_4 = 19,548.15$$

∴ At the beginning of 2013

Car is worth £19,548.15

Section 2 - Recurrence Relations.Additional Example 2.1.2

When a particular drug is given to a patient, 40% of it disappears from the body after each hour. 100mg of the drug is given to the patient at the start of treatment and 75mg after each hour.

- a) Write a recurrence relation for this situation.
 b) How much drug will be in the patient's body after 3 hours?

$$a) \quad U_{n+1} = 0.6U_n + 75 \quad U_0 = 100.$$

$$b) \quad U_1 = 0.6 \times 100 + 75$$

$$U_1 = 135$$

$$U_2 = 0.6 \times 135 + 75$$

$$U_2 = 156$$

$$U_3 = 0.6 \times 156 + 75$$

$$U_3 = 168.6$$

Ans 3 hrs

168.6mg

Example 4: Bill puts lottery winnings of £120000 in a bank account which pays 5% interest p/a. After a year, he decides to spend £20000 per year from the money in the account.

a) Find a recurrence relation to describe the amount of money left each year.

b) How much money will there be in the account after five years?

c) After how many years will Bill's money run out?


The Straight Line

Revision from National 5

The graph of $y = mx + c$ is a **straight line**, where m is the gradient and c is the y-intercept.

Gradient is a measure of the steepness of a line. The gradient of the line joining points A (x_1, y_1) and B (x_2, y_2) is given by:

$$m_{AB} = \frac{y_2 - y_1}{x_2 - x_1}$$



Example 1: Find:

a) the gradient and y-intercept of the line

$$y = 2x + 5$$

$$m = 2, c = 5$$

c) the gradient of the line joining P $(-2, 4)$ and Q $(3, -1)$

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m = \frac{4 - (-1)}{-2 - 3}$$

$$m = \frac{5}{-5}$$

$$m = -1$$

b) the equation of the line with gradient -4 and y-intercept $(0, -2)$

$$y = -4x - 2$$

d) the gradient of the line $3y + 4x - 11 = 0$

$$y = mx + c$$

$$3y = -4x + 11$$

$$y = -\frac{4}{3}x + \frac{11}{3}$$


$$m = -\frac{4}{3}$$

Equation of a Straight Line: $y - b = m(x - a)$

Points A (a, b) and P (x, y) both lie on a straight line.

The gradient of the line $m = \frac{y-b}{x-a}$. Rearranging this gives:

$y - b = m(x - a)$



NOTE: when you are asked to find the equation of a straight line, you should always expand the brackets and simplify as far as you can. If the question doesn't ask for a specific form then just leave it in any **simplified** form you want.

Example 2: Find the equations of the lines:

a) through (4, 5) with $m = 2$

$y - b = m(x - a)$
 $y - 5 = 2(x - 4)$
 $y - 5 = 2x - 8$
 $y = 2x - 3$

b) joining (-1, -2) and (3, 10)

x_1, y_1, x_2, y_2
 $m = \frac{y_2 - y_1}{x_2 - x_1}$
 $m = \frac{10 - (-2)}{3 - (-1)} = \frac{12}{4} = \underline{\underline{3}}$

c) parallel to the line $x - 2y + 4 = 0$ and passing through the point (2, -3)

$2y = x + 4$
 $y = \frac{1}{2}x + 2$ $m = \frac{1}{2}$
 $y - b = m(x - a)$
 $y + 3 = \frac{1}{2}(x - 2)$
 $y + 3 = \frac{1}{2}x - 1$
 $y = \frac{1}{2}x - 4$

$y - b = m(x - a)$
 $y - (-2) = 3(x - (-1))$
 $y + 2 = 3x + 3$
 $y = 3x + 1$


Equation of a Straight Line

$$y = mx + c \quad \text{AND} \quad Ax + By + C = 0$$

Example 3: Find the equation of the line through $(-5, -1)$ with $m = -\frac{2}{3}$, giving your answer in the form $Ax + By + C = 0$.

$$\begin{aligned} y - b &= m(x - a) \\ y - (-1) &= -\frac{2}{3}(x - (-5)) \\ y + 1 &= -\frac{2}{3}(x + 5) \\ 3y + 3 &= -2(x + 5) \\ 3y + 3 &= -2x - 10 \\ \underline{2x + 3y + 13} &= 0 \end{aligned}$$

Example 4: Sketch the line $5x - 2y - 24 = 0$ by finding the points where it crosses the x - and y -axes.




The Angle with the x-axis

The gradient of a line can also be described as the angle it makes with the **positive** direction of the x-axis.

As the y-difference is **OPPOSITE** the angle and the x-difference is **ADJACENT** to it, we get:

$$m_{AB} = \tan \theta$$

(where θ is measured **CLOCKWISE** from the x-axis)



Example 5: Find the angle made with the positive direction of the x-axis and the lines:

a) $y = x - 1$

Handwritten:
 $m = 1$
 $\tan \theta = m$
 $\tan \theta = 1$
 $\theta = \tan^{-1}(1)$
 $\theta = 45^\circ$ (R.A.)
 $\theta = 45^\circ$

b) $y = 5 - \sqrt{3}x$

Handwritten:
 $m = -\sqrt{3}$
 $\tan \theta = -\sqrt{3}$
 $\theta = \tan^{-1}(\sqrt{3})$
 $\theta = 60^\circ$ (R.A.)
 $\theta = 180 - 60$
 $= 120^\circ$

c) joining the points (3, -2) and (7, 4)

Handwritten:
 $m = \frac{y_2 - y_1}{x_2 - x_1}$
 $m = \frac{4 - (-2)}{7 - 3}$
 $m = \frac{6}{4} = \frac{3}{2}$
 $m = \tan \theta$
 $\tan \theta = \frac{3}{2}$
 $\theta = \tan^{-1}(\frac{3}{2})$
 $\theta = 56.3^\circ$ (R.A.)
 $\theta = 56.3^\circ$

Gradients of straight lines can be summarised as follows:

- a) lines sloping **up** from left to right have **positive** gradients and make **acute** angles with the positive direction of the x-axis
- b) lines sloping **down** from left to right have **negative** gradients and make **obtuse** angles with the positive direction of the x-axis
- c) lines with **equal** gradients are parallel
- d) **horizontal** lines (parallel to the x-axis) have gradient **zero** and equation $y = a$
- e) **vertical** lines (parallel to the y-axis) have gradient **undefined** and equation $x = b$

Collinearity

If three (or more) points lie on the same line, they are said to be collinear.

Example 6: Prove that the points D (-1, 5), E (0, 2) and F (4, -10) are collinear.

$$\begin{aligned}
 M_{DE} &= \frac{y_2 - y_1}{x_2 - x_1} \\
 &= \frac{5 - 2}{-1 - 0} \\
 &= \frac{3}{-1} \\
 &= -3.
 \end{aligned}$$

$$\begin{aligned}
 M_{EF} &= \frac{-10 - 2}{4 - 0} \\
 &= \frac{-12}{4} \\
 &= -3
 \end{aligned}$$

\therefore Since $M_{DE} = M_{EF}$
& E is a common point.
D, E & F are collinear.

Perpendicular Lines

If two lines are perpendicular to each other (i.e. they meet at 90°), then:

$$m_1 \times m_2 = -1$$

Example 7: Show whether these pairs of lines are perpendicular:

a) $x + y + 5 = 0$ $y = -x - 5$
 $x - y - 7 = 0$ $y = x - 7$

$$\begin{aligned}
 M_1 &= -1 \\
 M_2 &= 1 \\
 M_1 \times M_2 &= -1 \times 1 \\
 &= -1 \\
 \therefore \text{Since } M_1 M_2 &= -1 \\
 \text{Lines are perpendicular.}
 \end{aligned}$$

b) $2x - 3y = 5$ $M = \frac{2}{3}$
 $3x = 2y + 9$ $M = \frac{3}{2}$

$$\begin{aligned}
 M_1 &= \frac{2}{3} \\
 M_2 &= \frac{3}{2} \\
 \frac{2}{3} \times \frac{3}{2} &= 1 \\
 \therefore \text{Since } M_1 M_2 &\neq -1 \\
 \text{Lines are not} \\
 \text{perpendicular.}
 \end{aligned}$$

c) $y = 2x - 5$ $M = 2$
 $6y = 10 - 3x$ $M = -\frac{1}{2}$

$$\begin{aligned}
 M_1 &= 2 \\
 M_2 &= -\frac{1}{2} \\
 2 \times (-\frac{1}{2}) &= -1 \\
 \therefore \text{Since } M_1 M_2 &= -1 \\
 \text{Lines are} \\
 \text{perpendicular.}
 \end{aligned}$$

When asked to find the gradient of a line perpendicular to another, follow these steps:

1. Find the gradient of the given line
2. Flip it upside down
3. Change the sign (e.g. negative to positive)

Example 8: Find the gradients of the lines perpendicular to:

a) the line $y = 3x - 12$

$$M_1 = +3$$

$$M_2 = -\frac{1}{3}$$

b) a line with gradient = -1.5

$$M_1 = -\frac{3}{2}$$

$$M_2 = \frac{2}{3}$$

c) the line $2y + 5x = 0$

$$2y = -5x$$

$$y = -\frac{5}{2}x$$

$$M_1 = -\frac{5}{2}$$

$$M_2 = \frac{2}{5}$$

Example 9: Line L has equation $x + 4y + 2 = 0$. Find the equation of the line perpendicular to L which passes through the point $(-2, 5)$.

$$x + 4y + 2 = 0$$

$$4y = -x - 2$$

$$y = -\frac{1}{4}x - \frac{1}{2}$$

$$M_1 = -\frac{1}{4}$$

$$M_2 = 4 \quad (-2, 5)$$

$$y - b = m(x - a)$$

$$y - 5 = 4(x + 2)$$

$$y - 5 = 4x + 8$$

$$y = 4x + 13$$


Midpoints and Perpendicular Bisectors

The **midpoint** of a line lies exactly halfway along it. To find the coordinates of a midpoint, find halfway between the x - and y - coordinates of the points at each end of the line (see diagram).

The x - coordinate of M is halfway between -2 and 8, and its y - coordinate is halfway between 5 and -3.

In general, if M is the midpoint of A (x₁, y₁) and B (x₂, y₂):

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$



The **perpendicular bisector** of a line passes through its midpoint at 90°.

Example 10: Find the perpendicular bisector of the line joining F (-4, 2) and G (6, 8).

Handwritten calculations for Example 10:

$$M = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m = \frac{8 - 2}{6 - (-4)}$$

$$m = \frac{6}{10} = \frac{3}{5}$$

Gradient of FG is $\frac{3}{5}$.

Perpendicular gradient $m_{\perp} = -\frac{5}{3}$.

Midpoint $M = \left(\frac{-4+6}{2}, \frac{2+8}{2} \right) = (1, 5)$.

Equation of perpendicular bisector: $y - 5 = -\frac{5}{3}(x - 1)$.

To find the equation of a perpendicular bisector:

- ✓ Find the gradient of the line joining the given points
- ✓ Find the perpendicular gradient (flip and make negative)
- ✓ Find the coordinates of the midpoint
- Substitute into $y - b = m(x - a)$

Handwritten solution for the equation of the perpendicular bisector:

$$y - b = m(x - a)$$

$$y - 5 = -\frac{5}{3}(x - 1) \times 3$$


$$3y - 15 = -5(x - 1)$$

$$3y - 15 = -5x + 5$$

$$3y + 5x - 20 = 0$$


**Lines Inside Triangles:
Medians, Altitudes & Perpendicular Bisectors**

In a triangle, a line joining a corner to the **midpoint** of the opposite side is called a **median**.




The medians are **concurrent** (i.e. meet at the same point) at the **centroid**, which divides each median in the ratio 2:1. The median always divides the area of a triangle in half. A solid triangle of uniform density will balance on the centroid.

A line through a corner which is **perpendicular** to the opposite side is called an **altitude**.



The altitudes are concurrent at the **orthocentre**. The orthocentre isn't always located inside the triangle e.g. if the triangle is obtuse.

A line at 90° to the midpoint is called a **perpendicular bisector**.



The perpendicular bisectors are concurrent at the **circumcentre**. The circumcentre is the centre of the circle touched by the vertices of the triangle.

For all triangles, the centroid, orthocentre and circumcentre are **collinear**.

Lesson Starter - 5A1 - Thu 23/8/18


1) A & B are the points (4, 6) and (10, 8) respectively.

Find the equation of the perpendicular bisector of AB

$$M_{AB} = \left(\frac{4+10}{2}, \frac{6+8}{2} \right) \\ = (7, 7)$$

2) Find the length of the line CD

$$m_{AB} = \frac{y_2 - y_1}{x_2 - x_1} \\ = \frac{8 - 6}{10 - 4} \\ = \frac{2}{6} = \frac{1}{3} \\ m_{\perp} = -3$$




$$y - b = m(x - a) \\ y - 7 = -3(x - 7) \\ y - 7 = -3x + 21 \\ y = -3x + 28$$

$$CO^2 = 4^2 + 3^2 \\ CO^2 = 16 + 9 \\ CO^2 = 25 \\ CO = \sqrt{25} \\ = 5$$

Example 11: A triangle has vertices P (0, 2), Q (4, 4) and R (8, -6).

a) Find the equation of the median through P.

$$\begin{aligned}
 M_{QR} &= \left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2} \right) \\
 &= \left(\frac{4+8}{2}, \frac{4+(-6)}{2} \right) \\
 &= (6, -1) \\
 m_{PT} &= \frac{-1-2}{6-0} \\
 &= \frac{-3}{6} \\
 &= -\frac{1}{2} \\
 y-b &= m(x-a) \\
 y-2 &= -\frac{1}{2}(x-0) \\
 y-2 &= -\frac{1}{2}x \\
 \underline{y} &= \underline{-\frac{1}{2}x+2}
 \end{aligned}$$




- To find the equation of a median:
- Find the midpoint of the side opposite the given point
 - Find the gradient of the line joining the given point and the midpoint
 - Substitute into $y - b = m(x - a)$

b) Find the equation of the altitude through R.

$$\begin{aligned}
 m_{PQ} &= \frac{y_2 - y_1}{x_2 - x_1} & m_{\perp PQ} &= -2 \\
 &= \frac{4 - 2}{4 - 0} & y - b &= m(x - a) \\
 &= \frac{2}{4} & y - (-6) &= -2(x - 8) \\
 &= \frac{1}{2} & y + 6 &= -2x + 16 \\
 & & \underline{y} &= \underline{-2x + 10}
 \end{aligned}$$

- To find the equation of an altitude:
- Find the gradient of the side opposite the given point
 - Find the perpendicular gradient (flip and make negative)
 - Substitute into $y - b = m(x - a)$



$$\begin{aligned}
 y - b &= m(x - a) \\
 y - 1 &= -1(x - 3) \\
 y &= -x + 3 + 1 \\
 y &= -x + 4 \\
 \underline{\underline{y = -x + 4}}
 \end{aligned}$$

$$\begin{aligned}
 M_{AH} &= \frac{-6 - (-7)}{5 - 4} = \frac{-1}{1} \\
 &= -1 \\
 &= 1 \\
 \underline{\underline{M_{\perp} = -1}}
 \end{aligned}$$


Distance between Two Points

The distance between any two points A (x_1, y_1) and B (x_2, y_2) can be found easily by Pythagoras' Theorem.

If d is the distance between A and B, then:

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$d^2 = (x_2 - x_1)^2 + (y_2 - y_1)^2$$



Example 12: Calculate the distance between:

a) A $(-4, 4)$ and B $(2, -4)$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$d = \sqrt{(2 - (-4))^2 + (-4 - 4)^2}$$

$$d = \sqrt{36 + 64}$$

$$d = \sqrt{100}$$

$$d = 10$$

b) X $(11, 2)$ and Y $(-2, -5)$

$$d = \sqrt{(-2 - 11)^2 + (-5 - 2)^2}$$

$$d = \sqrt{169 + 49}$$

$$d = \sqrt{218}$$

Example 13: A is the point $(2, -1)$, B is $(5, -2)$ and C is $(7, 4)$. Show that $BC = 2AB$.

$$BC$$

$$d = \sqrt{(y_2 - y_1)^2 + (x_2 - x_1)^2}$$

$$= \sqrt{(-2 - 4)^2 + (5 - 7)^2}$$

$$= \sqrt{36 + 4}$$

$$= \sqrt{40} = 2\sqrt{10}$$

$2AB$

$$d = 2\sqrt{(1 - (-1))^2 + (2 - 5)^2}$$

$$d = 2\sqrt{1 + 9}$$

$$d = 2\sqrt{10}$$

\therefore Since $BC = 2\sqrt{10}$ & $2AB = 2\sqrt{10}$

$$BC = 2AB$$

Past Paper Example 1: The vertices of triangle ABC are A(7, 9), B(-3, -1) and C(5, -5) as shown: The broken line represents the perpendicular bisector of BC

a) Show that the equation of the perpendicular bisector of BC is $y = 2x - 5$

$$m_{BC} = \frac{-5 + 1}{5 + 3} = \frac{-4}{8} = -\frac{1}{2}$$

$$M_{BC} = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right) = \left(\frac{-3 + 5}{2}, \frac{-1 + (-5)}{2} \right) = (1, -3)$$

$$y - b = m(x - a)$$

$$y - (-3) = 2(x - 1)$$

$$y + 3 = 2x - 2$$

$$y = 2x - 5$$

b) Find the equation of the median from C

$$M_{AB} = \left(\frac{-3 + 7}{2}, \frac{-1 + 9}{2} \right) = (2, 4)$$

$$m_{AC} = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-5 - 4}{5 - 2} = \frac{-9}{3} = -3$$

$$y - b = m(x - a)$$

$$y + 5 = -3(x - 5)$$

$$y = -3x + 10$$

c) Find the co-ordinates of the point of intersection of the perpendicular bisector of BC and the median from C.

$$y = -3x + 10$$

$$y = 2x - 5$$

$$y = y$$

$$-3x + 10 = 2x - 5$$

$$-5x = -15$$

$$x = 3$$

When $x = 3$


$$y = 2(3) - 5$$

$$y = 1 \therefore \text{PoC } (3, 1)$$

Past Paper Example 2:

The line GH makes an angle of 30° with the y-axis as shown in the diagram opposite.

What is the gradient of GH?




$$\begin{aligned} m &= \tan \theta \\ m &= \tan 60^\circ \\ m &= \sqrt{3} \\ m &= 1.7 \end{aligned}$$

Vectors

Revision from National 5

A measurement which only describes the magnitude (i.e. size) of the object is called a **scalar** quantity, e.g. Glasgow is 11 miles from Airdrie. A **vector** is a quantity with **magnitude and direction**, e.g. Glasgow is 11 miles from Airdrie on a bearing of 270° .


The position of a point in 3-D space can be described if we add a third coordinate to indicate height.



Example 1: OABC DEFG is a cuboid, where F is the point (5, 4, 3). Write down the coordinates of the points:

- a) A
- b) D
- c) G
- d) M, the centre of face ABFE


The rules of vectors can be used in either 2 or 3 dimensions:



Directed line segment \overrightarrow{OA}

Position vector \underline{a}

Components $\begin{pmatrix} 3 \\ 4 \end{pmatrix}$



Directed line segment \overrightarrow{OB}

Position vector \underline{b}

Components $\begin{pmatrix} 2 \\ 5 \\ 4 \end{pmatrix}$

The **magnitude** of a vector is its length, which can be determined by Pythagoras' Theorem. The magnitude of \underline{a} is written as $|\underline{a}|$.

Example 2: Determine $|\underline{a}|$ and $|\underline{b}|$ in the examples above.

If $\underline{u} = \begin{pmatrix} a \\ b \end{pmatrix}$, then $|\underline{u}| = \sqrt{a^2 + b^2}$

If $\underline{u} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$, then $|\underline{u}| = \sqrt{a^2 + b^2 + c^2}$

Addition of Vectors

Two (or more) vectors can be added together to produce a **resultant vector**.

In general:

$$\vec{AB} + \vec{BC} = \vec{AC}$$

and


$$\text{If } \underline{u} = \begin{pmatrix} a \\ b \\ c \end{pmatrix} \text{ and } \underline{v} = \begin{pmatrix} d \\ e \\ f \end{pmatrix}, \text{ then } \underline{u} + \underline{v} = \begin{pmatrix} a + d \\ b + e \\ c + f \end{pmatrix}$$

Example 3: Find $\underline{p} + \underline{q}$ when $\underline{p} = \begin{pmatrix} 2 \\ 3 \\ -1 \end{pmatrix}$ and $\underline{q} = \begin{pmatrix} 7 \\ -7 \\ 4 \end{pmatrix}$.

Example 4: Find values of x and y such that

$$\begin{pmatrix} x \\ 4 \end{pmatrix} + \begin{pmatrix} 12 \\ y \end{pmatrix} = \begin{pmatrix} 9 \\ -2 \end{pmatrix}$$


Subtraction of Vectors



Subtraction of vectors can be considered as going along the second vector in the **wrong direction**.

$$\text{If } \underline{u} = \begin{pmatrix} a \\ b \\ c \end{pmatrix} \text{ and } \underline{v} = \begin{pmatrix} d \\ e \\ f \end{pmatrix}, \text{ then } \underline{u} - \underline{v} = \begin{pmatrix} a - d \\ b - e \\ c - f \end{pmatrix}$$

Multiplication by a Scalar Quantity



If we go along \underline{a} twice, the resultant vector is $\underline{a} + \underline{a} = 2\underline{a}$. As we have not changed direction, it follows that $2\underline{a}$ must be **parallel** to \underline{a} .

$$\text{If } \underline{u} = \begin{pmatrix} a \\ b \\ c \end{pmatrix} \text{ then } k\underline{u} = \begin{pmatrix} ka \\ kb \\ kc \end{pmatrix}$$

If $\underline{v} = k\underline{u}$, then \underline{u} and \underline{v} are parallel


Example 5: If $\underline{b} = \begin{pmatrix} 4 \\ 0 \\ -2 \end{pmatrix}$ and $\underline{c} = \begin{pmatrix} -3 \\ -5 \\ 5 \end{pmatrix}$, find:

a) $3\underline{b}$

b) $2\underline{b} + \underline{c}$

c) $\underline{c} - \frac{1}{2}\underline{b}$

Position Vectors



Consider the vector \vec{AB} in the diagram opposite. \vec{AB} is the resultant vector of going along \underline{a} in the opposite direction, followed by \underline{b} in the correct direction.

So, $\vec{AB} = -\underline{a} + \underline{b}$, i.e.:

$$\vec{AB} = \underline{b} - \underline{a}$$

Example 6: L is the point (4, -7, 2), M is the point (-5, -3, -1).

Find the components of \vec{LM} .

Handwritten blue notes:


$$\vec{LM} = \underline{m} - \underline{l}$$

$$= \begin{pmatrix} -5 \\ -3 \\ -1 \end{pmatrix} - \begin{pmatrix} 4 \\ -7 \\ 2 \end{pmatrix}$$

$$= \begin{pmatrix} -9 \\ 4 \\ -3 \end{pmatrix}$$

Example 7: P is the point (3, 7, -1). \vec{PQ} has components $\begin{pmatrix} -4 \\ 9 \\ -3 \end{pmatrix}$.

Find the coordinates of Q.



$$\vec{PQ} = \begin{pmatrix} -4 \\ 9 \\ -3 \end{pmatrix}$$

$$\vec{PQ} = \underline{Q} - \underline{P}$$

$$\begin{pmatrix} -4 \\ 9 \\ -3 \end{pmatrix} = \underline{Q} - \underline{P}$$

$$\begin{pmatrix} -4 \\ 9 \\ -3 \end{pmatrix} = \underline{Q} - \begin{pmatrix} 3 \\ 7 \\ -1 \end{pmatrix}$$

$$\underline{Q} = \begin{pmatrix} -4 \\ 9 \\ -3 \end{pmatrix} + \begin{pmatrix} 3 \\ 7 \\ -1 \end{pmatrix}$$

$$\underline{Q} = \begin{pmatrix} -1 \\ 16 \\ -4 \end{pmatrix}$$

$$\underline{Q} = \underline{(-7, 2, -2)}$$

C: x13G
Q1 #2

Unit Vectors

A unit vector is a vector with magnitude = 1.

Example 8: Find the components of the unit vector parallel to $\underline{h} = \begin{pmatrix} 2 \\ -3 \\ 6 \end{pmatrix}$

$$\begin{aligned} |\underline{h}| &= \sqrt{2^2 + (-3)^2 + 6^2} \\ &= \sqrt{4 + 9 + 36} \\ &= \sqrt{49} \\ &= \underline{7} \end{aligned}$$

$$\underline{u} = \begin{pmatrix} 2/7 \\ -3/7 \\ 6/7 \end{pmatrix}$$

To find the components of a unit vector:


- Find the magnitude of the given vector
- Divide components by the magnitude

Collinearity

Example 9: Points F, G and H have coordinates (6, 1, 5), G (4, 4, 4), and (-2, 13, 1) respectively. Show that F, G and H are collinear, and find the ratio in which G divides FH.

$$\begin{aligned}\vec{FG} &= \vec{g} - \vec{f} \\ &= \begin{pmatrix} 4 \\ 4 \\ 4 \end{pmatrix} - \begin{pmatrix} 6 \\ 1 \\ 5 \end{pmatrix} \\ &= \begin{pmatrix} -2 \\ 3 \\ -1 \end{pmatrix}\end{aligned}\qquad \begin{aligned}\vec{GH} &= \vec{h} - \vec{g} \\ &= \begin{pmatrix} -2 \\ 13 \\ 1 \end{pmatrix} - \begin{pmatrix} 4 \\ 4 \\ 4 \end{pmatrix}\end{aligned}$$

\therefore Since $\vec{GH} = 3\vec{FG}$ & G is a common point,
F, G & H are collinear & parallel.





Example 10: The points $P(6, 1, -3)$, $Q(8, -3, 1)$ and $R(9, -5, 3)$ are collinear.
Find the ratio in which Q divides PR

The Section Formula

P divides \overline{AB} in the ratio 2:3. By examining the diagram, we can find a formula for \underline{p} (i.e. \overline{OP}).

$$\overline{OP} = \overline{OA} + \overline{AP}$$



In general, if P divides AB in the ratio m:n, then:

$$\underline{p} = \frac{1}{n+m} (n\underline{a} + m\underline{b})$$

Example 11: A is the point (3, -1, 2) and B is the point (7, -5, 14). Find the coordinates of P such that P divides AB in the ratio 1:3.

Vectors in 3D can also be described in terms of the three unit vectors $\underline{i} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$, $\underline{j} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$, and $\underline{k} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$,
which are parallel to the x, y, and z axes respectively.

Example 12: $\underline{u} = 3\underline{i} + 2\underline{j} - 6\underline{k}$, $\underline{v} = -\underline{i} + 5\underline{j}$.

a) Express $\underline{u} + \underline{v}$ in component form

b) Find $|\underline{u} + \underline{v}|$


The Scalar Product (Angle Form)

The **scalar product** is the result of a type of multiplication of two vectors to give a scalar quantity. (i.e. a number with no directional component)

For vectors \underline{a} and \underline{b} , the scalar product (or **dot product**) is given as:


$$\underline{a} \cdot \underline{b} = |\underline{a}| |\underline{b}| \cos \theta$$

- Note:
- \underline{a} and \underline{b} point away from the vertex
 - $0 \leq \theta \leq 180^\circ$




Example 13: Find the scalar product in each case below, where $|\underline{a}| = 6$ and $|\underline{b}| = 10$.


a)



b)



c)



Two vertical lines for writing answers to parts b) and c).

The Scalar Product (Component Form)

We can use the formula below to find the scalar product when we have been given the component forms of the two vectors but not the angle in between them.


$$\text{If } \underline{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \text{ and } \underline{b} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}, \text{ then } \underline{a} \cdot \underline{b} = a_1b_1 + a_2b_2 + a_3b_3$$

Example 14: $\underline{a} = i + 2j + 2k$, and $\underline{b} = 2i + 3j - 6k$. Evaluate $\underline{a} \cdot \underline{b}$

Example 15: A is the point (1, 2, 3), B is the point (6, 5, 4) and C is the point (-1, -2, -6).
Evaluate $\overline{AB} \cdot \overline{BC}$

Perpendicular Vectors

A special case of the scalar product occurs when we have perpendicular vectors i.e. when $\theta = 90^\circ$:



$$\begin{aligned} a \cdot b &= |a| |b| \cos 90^\circ \\ &= |a| |b| \times 0 \\ &= 0 \end{aligned}$$

If $a \cdot b = 0$, then a and b are perpendicular

Example 16: P, Q and R are the points (1, 1, 2), (-1, -1, 0) and (3, -4, -1) respectively. Find the components of \overline{QP} and \overline{QR} , and hence show that the vectors are perpendicular.

$$\underline{a}\underline{i} + \underline{b}\underline{j} + \underline{k} \perp 3\underline{i} - 4\underline{j} + 6\underline{k}$$

$$-5\underline{i} + 2\underline{k} + 4\underline{j}$$

$$\underline{u} = \begin{pmatrix} a \\ b \\ 1 \end{pmatrix}$$

$$\underline{v} = \begin{pmatrix} 3 \\ -4 \\ 6 \end{pmatrix}$$

$$\underline{w} = \begin{pmatrix} -5 \\ 4 \\ 2 \end{pmatrix}$$

$$\underline{u} \cdot \underline{v} = 3a - 4b + 6$$

perpendicular vectors happen
when $\underline{u} \cdot \underline{v} = 0$.

$$3a - 4b + 6 = 0.$$

$$\underline{u} \cdot \underline{w} = -5a + 2b + 4$$

$$-5a + 2b + 4 = 0$$

$$3a - 4b = -6$$

$$-5a + 2b = -4 \quad (\times 2)$$

$$\begin{array}{r} 3a - 4b = -6 \\ -10a + 4b = -8 \end{array} \quad \underline{\text{Add}}.$$

$$\begin{array}{r} -7a = -14 \\ a = 2 \end{array}$$

$$\text{When } a=2, \quad 3(2) - 4b + 6 = 0$$

$$6 - 4b + 6 = 0$$

$$4b = 12$$

$$b = 3.$$

$$\therefore a = 2$$

$$b = 3.$$

The Angle Between Two Vectors

We can rearrange the angle form of the scalar product to give $\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}| |\mathbf{b}|}$.

More specifically:

$$\cos ABC = \frac{\overline{BA} \cdot \overline{BC}}{|\overline{BA}| |\overline{BC}|}$$

If the question gives you three points, you MUST find the components of the vectors pointing AWAY from the vertex first!

Example 17: $\mathbf{a} = i + 2j + 2k$ and $\mathbf{b} = 2i + 3j - 6k$. Find the angle between \mathbf{a} and \mathbf{b} .

$$\mathbf{a} = \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$$

$$\mathbf{b} = \begin{pmatrix} 2 \\ 3 \\ -6 \end{pmatrix}$$

Ⓢ
NANA

$$\cos ABC = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}| |\mathbf{b}|}$$

$$\cos \theta = \frac{-4}{3 \times 7}$$

$$= \frac{-4}{21}$$

$$\theta = \cos^{-1} \left(\frac{-4}{21} \right)$$

$$\theta = 79.01^\circ \text{ (R.A.)}$$

$$\theta = 180 - 79.01^\circ$$

$$\theta = \underline{100.9^\circ}$$

$$|\mathbf{a}|$$

$$= \sqrt{1^2 + 2^2 + 2^2}$$

$$= \sqrt{1 + 4 + 4}$$

$$= \sqrt{9} = 3$$


$$|\mathbf{b}| = \sqrt{2^2 + 3^2 + 6^2}$$

$$= \sqrt{4 + 9 + 36}$$

$$= \sqrt{49}$$

$$= 7$$

Example 18: A is the point (1, 2, 3), B (6, 5, 4), and C (-1, -2, -6). Calculate $\angle ABC$.



$$\begin{aligned} \vec{BA} &= \underline{a} - \underline{b} \\ &= \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} - \begin{pmatrix} 6 \\ 5 \\ 4 \end{pmatrix} \\ &= \begin{pmatrix} -5 \\ -3 \\ -1 \end{pmatrix} \cdot 19 \end{aligned}$$

$$\begin{aligned} \vec{BC} &= \underline{c} - \underline{b} \\ &= \begin{pmatrix} -1 \\ -2 \\ -6 \end{pmatrix} - \begin{pmatrix} 6 \\ 5 \\ 4 \end{pmatrix} \\ &= \begin{pmatrix} -7 \\ -7 \\ -10 \end{pmatrix} \cdot 15 \end{aligned}$$

$$\cos \theta = \frac{a \cdot b}{|a| |b|}$$

$$\cos \theta = \frac{66}{\sqrt{35} \times \sqrt{198}}$$

$$\cos \theta = 0.79$$

$$\begin{aligned} \theta &= \cos^{-1}(0.79) \\ &= \underline{37.8^\circ} \end{aligned}$$

$$\begin{aligned} \theta &= \cos^{-1} \left(\frac{66}{\sqrt{35} \times \sqrt{198}} \right) \\ \theta &= \underline{37.8^\circ} \end{aligned}$$


$$\begin{aligned} & \underline{a \cdot b} \\ &= -5 \times (-7) \\ &+ (-3) \times (-7) \\ &+ (-1) \times (-10) \\ &= 35 + 21 + 10 \\ &= \underline{66} \checkmark \\ & \underline{|a|} \\ &= \sqrt{-5^2 + (-3)^2 + (-1)^2} \\ &= \sqrt{25 + 9 + 1} \\ &= \underline{\sqrt{35}} \checkmark \\ & \underline{|b|} \\ &= \sqrt{(-7)^2 + (-7)^2 + (-10)^2} \\ &= \underline{\sqrt{198}} \checkmark \end{aligned}$$

Other Uses of the Scalar Product

For vectors \underline{a} , \underline{b} , and \underline{c} :

$$\underline{a} \cdot \underline{b} = \underline{b} \cdot \underline{a}$$

$$\underline{a} \cdot (\underline{b} + \underline{c}) = \underline{a} \cdot \underline{b} + \underline{a} \cdot \underline{c}$$

Example 19: $|\underline{a}| = 5$ and $|\underline{b}| = 8$. Find $\underline{a} \cdot (\underline{a} + \underline{b})$ 

$$\begin{aligned} & \underline{a} \cdot \underline{a} + \underline{a} \cdot \underline{b} \\ & 25 + \frac{40}{\sqrt{2}} \\ & = \underline{\underline{53.28}} \end{aligned}$$

$$\begin{aligned} & \underline{a} \cdot \underline{a} \\ & |\underline{a}| \times |\underline{a}| \times \cos 0 \\ & 5 \times 5 \times \cos 0 \\ & = 5 \times 5 \times 1 \\ & = \underline{\underline{25}} \\ & \underline{a} \cdot \underline{b} \\ & |\underline{a}| \times |\underline{b}| \times \cos \theta \\ & 5 \times 8 \times \cos 45^\circ \\ & = 40 \times \frac{1}{\sqrt{2}} \\ & = \underline{\underline{\frac{40}{\sqrt{2}}}} \end{aligned}$$

Lesson Starter - 5B1 - Mon 17/9/18

1) If $f(x) = 2x - 16$. Find $f^{-1}(x)$

$$y = 2x - 16$$


$$y + 16 = 2x$$

$$x = \frac{y + 16}{2}$$

$$f^{-1}(x) = \frac{x + 16}{2} \checkmark$$

2) Shown below is the graph of $y = f(x)$

Sketch the graph of $y = 2f(x + 1)$




$$= \frac{1}{2}x + \frac{16}{2}$$

$$= \frac{1}{2}x + 8$$

$(-6, 0) \rightarrow$	$(-6, 0) \rightarrow$	$(-7, 0)$
$(-2, 4) \rightarrow$	$(-2, 8) \rightarrow$	$(-3, 8)$
$(0, 0) \rightarrow$	$(0, 0) \rightarrow$	$(-1, 0)$
$(2, -2) \rightarrow$	$(2, -4) \rightarrow$	$(1, -4)$
$(4, 0) \rightarrow$	$(4, 0) \rightarrow$	$(3, 0)$

Past Paper Example 1: The diagram shows a square-based pyramid of height 8 units. Square OABC has a side length of 6 units. The coordinates of A and D are (6, 0, 0) and (3, 3, 8). C lies on the y - axis.

- a) Write down the coordinates of B.
- b) Determine the components of \overrightarrow{DA} and \overrightarrow{DB} .
- c) Calculate the size of $\angle ADB$.




Past Paper Example 2:

a) Show that the points A (-7, -8, 1), T (3, 2, 5) and B (18, 17, 11) are collinear and state the ratio in which T divides AB.

$$\vec{AT} = t - a = \begin{pmatrix} 3 \\ 2 \\ 5 \end{pmatrix} - \begin{pmatrix} -7 \\ -8 \\ 1 \end{pmatrix} = \begin{pmatrix} 10 \\ 10 \\ 4 \end{pmatrix}$$

$$\vec{TB} = b - t = \begin{pmatrix} 18 \\ 17 \\ 11 \end{pmatrix} - \begin{pmatrix} 3 \\ 2 \\ 5 \end{pmatrix} = \begin{pmatrix} 15 \\ 15 \\ 6 \end{pmatrix}$$

\therefore Since $\vec{TB} = \frac{3}{2} \vec{AT}$ & T is a common point.
A, T & B are collinear & parallel.



$$1 : \frac{3}{2} \quad \times 2.$$

$$\underline{\underline{2 : 3}}$$

b) The point C lies on the x-axis.

If TB and TC are perpendicular find the coordinates of C.

$$\vec{TB} = \begin{pmatrix} 15 \\ 15 \\ 6 \end{pmatrix} \quad \vec{TC} = c - t = \begin{pmatrix} x \\ 0 \\ 0 \end{pmatrix} - \begin{pmatrix} 3 \\ 2 \\ 5 \end{pmatrix} = \begin{pmatrix} x-3 \\ -2 \\ -5 \end{pmatrix}$$

$\vec{TB} \cdot \vec{TC} = 0$


$$\begin{aligned} \vec{TB} \cdot \vec{TC} &= 15(x-3) + 15(-2) + 6(-5) \\ &= 15x - 45 - 30 - 30 \\ &= 15x - 105 \end{aligned}$$

$$\begin{aligned} 15x - 105 &= 0 \\ 15x &= 105 \\ x &= 7 \end{aligned}$$

C(7, 0, 0)

Past Paper Example 3: PQRSTU is a regular hexagon of side 2 units. \overrightarrow{PQ} , \overrightarrow{QR} and \overrightarrow{RS} represent the vectors \underline{a} , \underline{b} and \underline{c} respectively. Find the value of $\underline{a} \cdot (\underline{b} + \underline{c})$

$$\begin{aligned}
 &= \underline{a} \cdot \underline{b} + \underline{a} \cdot \underline{c} \\
 &= 2 + (-2) \\
 &= 0
 \end{aligned}$$



$$\begin{aligned}
 \underline{a} \cdot \underline{b} &= |\underline{a}| |\underline{b}| \cos 60 \\
 &= 2 \times 2 \times \cos 60 \\
 &= 4 \times \frac{1}{2} \\
 &= 2
 \end{aligned}$$


$$\begin{aligned}
 \underline{a} \cdot \underline{c} &= 2 \times 2 \times \cos 120 \\
 &= -2
 \end{aligned}$$

Calculus 1: Differentiation

In the chapter on straight lines, we saw that the gradient of a line is a measure of how quickly it increases (or decreases) at a constant rate.

This is easy to see for linear functions, but what about quadratic, cubic and higher functions? As these functions produce curved graphs, they do **not** increase or decrease at a constant rate.

For a function $f(x)$, the rate of change at any point on the function can be found by measuring the gradient of a **tangent** to the curve at that point.



The rate of change at any point of a function is called the **derived function** or the **derivative**.

Finding the rate of change of a function at a given point is part of a branch of maths known as **calculus**.

For function $f(x)$ or the graph $y = f(x)$, the derivative is written as:

$f'(x)$ ("f dash x")

OR

$\frac{dy}{dx}$ ("dy by dx")

Derivative = Rate of Change of the Function = Gradient of the Tangent to the Curve

The Derivative of $f(x) = ax^n$

Example 1: Find the derivative of $f(x) = x^2$

To find the derivative of a function:

1. Make sure it's written in the form $y = ax^n$
2. Multiply by the power
3. Decrease the power by one

Handwritten work for Example 1:

$$f(x) = x^2$$

$$f'(x) = \underline{\underline{2x}} \quad \frac{dy}{dx} = \underline{\underline{2x}}$$

Example 2: $f(x) = 2x^3$. Find $f'(x)$.

This means:

At $x = 1$, the gradient of the tangent to $2x^3 =$

At $x = -2$, the gradient of the tangent to $2x^3 =$

If $f(x) = ax^n$, then $f'(x) = nax^{n-1}$

The DE rivative DE creases the power!

To find the derivative of $f(x)$:

- $f(x)$ MUST be written in the form $f(x) = ax^n$
- Rewrite to eliminate fractions by using negative indices
- Rewrite to eliminate roots by using fractional indices

Example 3: Differentiate the following expressions:

a) $f(x) = 6x^{10}$

b) $h(p) = 3p^{-5}$

c) $y = -12x^{-3}$

d) $k = -9x^{-9}$

Revision from National 5

Example 4: Write with negative indices:

a) $\frac{2}{x^2}$
 $2x^{-2}$

b) $\frac{1}{4x^5}$
 $\frac{1}{4}x^{-5}$

c) $\frac{3}{5x}$
 $\frac{3}{5}x^{-1}$

Example 5: Write in index form:

a) \sqrt{x}
 $= x^{1/2}$

b) $\sqrt[3]{x^2}$
 $= x^{2/3}$

c) $\frac{2}{3\sqrt{x^7}}$
 $= \frac{2}{3}x^{-7/2}$

Handwritten work for Example 4b:

$$\frac{1}{4} \times \frac{1}{x^5}$$

$$\frac{1}{4} \times \frac{x^{-5}}{1}$$

$$\frac{1}{4} x^{-5}$$

Example 6: For each function, find the derivative.

a) $f(x) = \frac{4}{x^3}$

$$f(x) = 4x^{-3}$$

$$f'(x) = -12x^{-4}$$

$$f'(x) = \frac{-12}{x^4}$$

b) $g(x) = \frac{3}{4x^5}$

$$g(x) = \frac{3}{4}x^{-5}$$

$$g'(x) = \frac{-15}{4}x^{-6}$$

$$g'(x) = \frac{-15}{4x^6}$$

c) $p(x) = \frac{1}{\sqrt{x}} \quad (x > 0)$

$$p(x) = \frac{1}{x^{1/2}}$$

$$p(x) = x^{-1/2}$$

$$p'(x) = -\frac{1}{2}x^{-3/2}$$

$$= \frac{-1}{2x^{3/2}}$$

$$= \frac{-1}{2\sqrt{x^3}}$$

d) $y = 12x^5 + 3x^2 - 2x + 9$

$$\frac{dy}{dx} = 60x^4 + 6x - 2$$

e) $y = \frac{1}{3\sqrt{x}} \quad (x > 0)$

f) $y = (x^3 - 2)^2 \quad (x \geq 0)$

$$y = x^6 - 4x^3 + 4$$

$$\frac{dy}{dx} = 6x^5 - 12x^2$$

Example 7: Find the rate of change of each function:

$$\text{a) } f(x) = \frac{x^5 - 6x^3}{x^2}$$

$$f(x) = \frac{x^5}{x^2} - \frac{6x^3}{x^2}$$

$$f(x) = x^3 - 6x$$

$$f'(x) = 3x^2 - 6.$$

$$\text{b) } y = \frac{(x+3)^2}{x^{2/3}}$$

$$y = \frac{x^2 + 6x + 9}{x^{2/3}}$$

$$y = \frac{x^2}{x^{2/3}} + \frac{6x^1}{x^{2/3}} + \frac{9}{x^{2/3}}$$

$$y = x^{4/3} + 6x^{1/3} + 9x^{-2/3}$$

$$\frac{dy}{dx} = \frac{4}{3}x^{1/3} + \frac{6}{3}x^{-2/3} - 6x^{-5/3}$$

$$= \frac{4}{3}\sqrt[3]{x} + \frac{2}{\sqrt[3]{x^2}} - \frac{6}{3\sqrt[3]{x^5}}$$

$$1 - \frac{2}{3} = \frac{1}{3}$$

$$\text{c) } f(x) = \frac{x^5 - 3x}{2x^3}$$

$$f(x) = \frac{x^5}{2x^3} - \frac{3x}{2x^3}$$

$$f(x) = \frac{1}{2}x^2 - \frac{3}{2}x^{-2}$$

$$f'(x) = x + 3x^{-3}$$

$$= x + \frac{3}{x^3}$$

- Points to note:
- Number terms disappear (e.g. if $f(x) = 5$, $f'(x) = 0$)
 - x - terms leave their coefficient (e.g. if $f(x) = 135x$, $f'(x) = 135$)
 - Give your answer back in the same form as the question

Equation of a Tangent to a Curve

Example 8: Find the equation of the tangent to the curve $y = x^2 - 2x - 15$ when $x = 4$.


- To find the equation of a tangent to a curve:
- Find the point of contact (sub the value of x into the equation to find y)
 - Find $\frac{dy}{dx}$
 - Find m by substituting x into $\frac{dy}{dx}$
 - Use $y - b = m(x - a)$

Example 9:

a) Find the gradient of the tangent to the curve

$$y = x^3 - 2x^2 \text{ at the point where } x = \frac{7}{3}.$$


b) Find the other point on the curve where the tangent has the same gradient.



Example 10: Find the point of contact of the tangent to the curve with equation $y = x^2 + 7x + 3$ when the gradient of the tangent is 9.

Stationary Points and their Nature

Any point on a curve where the tangent is horizontal (i.e. the gradient or $\frac{dy}{dx} = 0$) is commonly known as a **stationary point**. There are four types of stationary point:



To locate the position of stationary points, we find the derivative, make it equal zero, and solve for x . To determine their type (or **nature**), we must use a **nature table**.

Example 11: Find the stationary points of the curve $y = 2x^3 - 12x^2 + 18x$ and determine their nature.

$$\frac{dy}{dx} = 6x^2 - 24x + 18.$$

SP's occur when $\frac{dy}{dx} = 0$.

$$6x^2 - 24x + 18 = 0.$$

$$6(x^2 - 4x + 3) = 0 \quad x = 3$$

$$6(x-3)(x-1) = 0$$

$$\underline{x=3}, \underline{x=1}$$

x	0	1	2	3	4
		→		→	
$\frac{dy}{dx}$	+	0	-	0	+
Shape		/	\	/	

When $x=3$
 $y = 2(3)^3 - 12(3)^2 + 18(3)$
 $y = 0$

When $x=1$
 $y = 2(1)^3 - 12(1)^2 + 18(1)$
 $y = 8$

When $x=0$

$$\frac{dy}{dx} = 6(0)^2 - 24(0) + 18$$

$$= 18$$

When $x=2$

$$\frac{dy}{dx} = 6(2)^2 - 24(2) + 18$$

$$= 24 - 48 + 18$$

$$= -6$$

When $x=4$

$$\frac{dy}{dx} = 6(4)^2 - 24(4) + 18$$

$$= 18$$

MaxTP @ (1, 8)

MinTP @ (3, 0)


Increasing & Decreasing Functions

if $\frac{dy}{dx} > 0$, then y is increasing

For any curve,

if $\frac{dy}{dx} < 0$, then y is decreasing

if $\frac{dy}{dx} = 0$, then y is stationary



If a function is **always** increasing (or decreasing), it is said to be **strictly** increasing (or decreasing).

Example 12: State whether the function $f(x) = x^3 - x^2 - 5x + 2$ is increasing, decreasing or stationary when:

a) $x = 0$

$$f'(x) = 3x^2 - 2x - 5$$

$$f'(0) = 3(0)^2 - 2(0) - 5 = -5$$

Decreasing.

b) $x = 1$

$$f'(1) = 3(1)^2 - 2(1) - 5 = 3 - 2 - 5 = -4$$

Decreasing.

c) $x = 2$

$$f'(2) = 3(2)^2 - 2(2) - 5 = 12 - 4 - 5 = 3$$

Increasing.

Example 13: Show algebraically that the function $f(x) = x^3 - 6x^2 + 12x - 5$ is never decreasing.

Example 14: Find the intervals in which the function $f(x) = 2x^3 - 6x^2 + 5$ is increasing and decreasing.

Curve Sketching

To accurately sketch and annotate the curve obtained from a function, we must consider:

1. x - and y - intercepts
2. Stationary points and their nature

Example 15: Sketch and annotate fully $y = x^3(4-x)$

Curve cuts x-axis when $y = 0$.

$$x^3(4-x) = 0$$

$$x^3 = 0, 4-x = 0$$

$$x = 0, x = 4$$

Curve cuts y-axis when $x = 0$

$$y = 0^3(4-0)$$

$$y = 0$$

$$y = x^3(4-x)$$

$$y = 4x^3 - x^4$$

$$\frac{dy}{dx} = 12x^2 - 4x^3$$

SPs occur when $\frac{dy}{dx} = 0$

$$12x^2 - 4x^3 = 0$$

$$4x^2(3-x) = 0$$

$$4x^2 = 0, 3-x = 0$$

$$x = 0, x = 3$$

When $x = 0$

$$y = 0^3(4-0)$$

$$y = 0$$

When $x = 3$

$$y = 3^3(4-3)$$

$$y = 27$$

Rising PE (0,0)
Max TP (3,27)

(4,0)
(0,0)

(0,0)

x	→ 0	→ 3	→
$\frac{dy}{dx}$	+	+	-
Shape	/ - / - \		

When $x = -1$

$$\frac{dy}{dx} = 12(-1)^2 - 4(-1)^3$$

$$= 12 + 4$$

$$= 16$$

When $x = 2$

$$\frac{dy}{dx} = 12(2)^2 - 4(2)^3$$

$$= 12(4) - 4(8)$$

$$= 48 - 32$$


$$= 16$$

When $x = 4$

$$\frac{dy}{dx} = 12(4)^2 - 4(4)^3$$

$$= 192 - 256$$


$$= -64$$



Closed Intervals


Sometimes, we may only be interested in a small section of the curve of a function. To find the maximum and minimum values of a function in a given interval, we find stationary points as normal, but we also need to consider the value of the function at the ends of the interval.

Example 16: Find the greatest and least values of $y = x^3 - 12x$ in the interval $-3 \leq x \leq 1$.



Note: In a closed interval. The maximum and minimum values of a function occur either at a Stationary Point within the interval or at the end point of the interval.


Graph of the Derived Function





From the graph of $y = f(x)$, we can obtain the graph of $y = f'(x)$ by considering its stationary points. On the graph of $y = f'(x)$, the y-coordinate comes from the derivative of $y = f(x)$.

1. Draw a set of axes directly under a copy of $y = f(x)$.
2. Locate the stationary points.
3. At SP's, $f'(x) = 0$, so the y coordinate of $f'(x) = 0$ on the new graph.
4. Where $f(x)$ is increasing, $f'(x)$ is **above** the x - axis.
5. Where $f(x)$ is decreasing, $f'(x)$ is **below** the x - axis.
6. Draw a smooth curve which fits this information.

Example 17: For the graphs below. Sketch the corresponding derived graphs of $y = f'(x)$








Differentiating $\sin x$ & $\cos x$

Let $f(x) = \sin x$ and $g(x) = \cos x$. The graphs of $y = f(x)$ and $y = g(x)$ are shown below, where the x -axis is measured in radians. Tangents to each curve have been drawn at the following points:

On $y = \sin x$, the tangent at $x = 0$ has $m = 1$, and the tangent at $x = \pi$ has $m = -1$.

On $y = \cos x$, the tangent at $x = \frac{\pi}{2}$ has $m = -1$, and the tangent at $x = \frac{3\pi}{2}$ has $m = 1$.

Draw the graphs of $y = f'(x)$ and $y = g'(x)$ below.



The graphs of the derived functions therefore show that:

If $y = \sin x$, $\frac{dy}{dx} = \cos x$

If $y = \cos x$, $\frac{dy}{dx} = -\sin x$.

Example 18: Find the derivative in each case:

a) $y = 4\sin x$

$$\frac{dy}{dx} = 4\cos x$$

b) $f(x) = 2\cos x$

$$f'(x) = -2\sin x$$

c) $g(x) = -\frac{1}{2}\cos x$

$$g'(x) = \frac{1}{2}\sin x$$

d) $h = -5\sin k$

$$\frac{dh}{dk} = -5\cos k$$

Past Paper Example 1: A curve has equation $y = x^4 - 4x^3 + 3$. Find the position and nature of its stationary points.

$$\frac{dy}{dx} = 4x^3 - 12x^2$$

SP's when $\frac{dy}{dx} = 0$.

$$4x^3 - 12x^2 = 0$$

$$4x^2(x - 3) = 0$$

$$4x^2 = 0, x - 3 = 0$$

$$x = 0, x = 3.$$

$$\text{When } x = 0, y = 0^4 - 4(0)^3 + 3 \quad (0, 3)$$

$$\text{When } x = 3, y = 3^4 - 4(3)^3 + 3 \quad (3, -24)$$

$$y = -24$$

$$\text{When } x = -1$$

$$\frac{dy}{dx} = 4(-1)^3 - 12(-1)^2$$

$$= -4 - 12$$

$$= -16$$

$$\text{When } x = 1$$

$$\frac{dy}{dx} = 4(1)^3 - 12(1)^2$$

$$= 4 - 12$$

$$= -8$$

$$\text{When } x = 4$$

$$\frac{dy}{dx} = 4(4)^3 - 12(4)^2$$

$$= 256 - 192$$

$$= 64$$

x	$\rightarrow 0$	$\rightarrow 3$	\rightarrow
$\frac{dy}{dx}$	$-$	0	$- 0 +$
Shape	$ $	$_ _ /$	

Local Max @ $(0, 3)$
Local Min @ $(3, -24)$

Past Paper Example 2: Find the equation of the two tangents to the curve $y = 2x^3 - 3x^2 - 12x + 20$ which are parallel to the line $48x - 2y = 5$.

$$2y = 48x + 5$$

$$y = 24x + \frac{5}{2} \quad m = 24$$

$$\frac{dy}{dx} = \frac{6x^2 - 6x - 12}{1}$$

$$6x^2 - 6x - 12 = 24$$

$$6x^2 - 6x - 36 = 0$$

$$6(x^2 - x - 6) = 0$$

$$6(x-3)(x+2) = 0$$

$$x = 3, x = -2$$

When $x = 3$, $y = 2(3)^3 - 3(3)^2 - 12(3) + 20$

$$y = 54 - 27 - 36 + 20$$

$$y = 11$$

(3, 11)

When $x = -2$, $y = 2(-2)^3 - 3(-2)^2 - 12(-2) + 20$

$$y = -16 - 12 + 24 + 20$$

$$y = 16$$

(-2, 16)

$$y - b = m(x - a) \quad \left| \quad y - 16 = 24(x + 2)$$

$$y - 11 = 24(x - 3) \quad \left| \quad y - 16 = 24x + 48$$

$$y - 11 = 24x - 72 \quad \left| \quad y = 24x + 64$$

$$\underline{y = 24x - 61}$$

Past Paper Example 3: A function is defined on the domain $0 \leq x \leq 3$ by $f(x) = x^3 - 2x^2 - 4x + 6$.

Determine the maximum and minimum values of f .

$$\text{When } x=0, y = 0^3 - 2(0)^2 - 4(0) + 6$$

$$y = 6$$

$(0, 6)$

$$\text{When } x=3, y = 3^3 - 2(3)^2 - 4(3) + 6$$

$$y = 27 - 18 - 12 + 6$$

$$y = 3$$

$(3, 3)$

$$\frac{dy}{dx} = 3x^2 - 4x - 4$$

$$\text{SP's when } \frac{dy}{dx} = 0$$

$$3x^2 - 4x - 4 = 0 \quad \begin{matrix} 3x+2 \\ x-2 \end{matrix}$$

$$(3x+2)(x-2) = 0$$

$$x = -\frac{2}{3}, x = 2$$

$$\text{When } x = -\frac{2}{3}, y = \left(-\frac{2}{3}\right)^3 - 2\left(-\frac{2}{3}\right)^2 - 4\left(-\frac{2}{3}\right) + 6$$

$$y = \frac{202}{27}$$

OUT OF RANGE

$\left(-\frac{2}{3}, 7.5\right)$

$$\text{When } x=2, y = 2^3 - 2(2)^2 - 4(2) + 6$$

$$y = 8 - 8 - 8 + 6$$

$$y = -2$$


$(2, -2)$

Max value = 6 when $x=0$

Min value = -2 when $x=2$.

The Circle

If we draw, suitable to relative axes, a circle, radius r , centred on the origin, then the distance from the centre of any point $P(x, y)$ could be determined to be $d = \sqrt{x^2 + y^2}$.



As the shape is a circle, then this distance is equal to the radius. It therefore follows that:

Since $r = \sqrt{x^2 + y^2}$, then $r^2 = x^2 + y^2$

Therefore,

The equation $x^2 + y^2 = r^2$ describes a circle with centre $(0, 0)$ and radius r

Example 1: Write down the centre and radius of each circle.

a) $x^2 + y^2 = 64$
 $(0, 0)$
 $r = 8$

b) $x^2 + y^2 = 361$
 $(0, 0)$
 $r = \sqrt{361}$
 $r = 19$

c) $x^2 + y^2 = \frac{3}{25}$
 $(0, 0)$
 $r = \sqrt{\frac{3}{25}}$ $r = \frac{\sqrt{3}}{5}$

Example 2: State where the points $(-2, 7)$, $(6, -8)$ and $(5, 9)$ lie in relation to the circle $x^2 + y^2 = 100$.

$(-2, 7)$
 $x^2 + y^2 = 100$
 $(-2)^2 + 7^2$
 $= 4 + 49$
 $= 53$
 \therefore Since $53 < 100$
 $(-2, 7)$ lies inside the circle.

$(6, -8)$
 $x^2 + y^2 = 100$
 $6^2 + (-8)^2$
 $= 36 + 64$
 $= 100$
 \therefore Since $100 = 100$
 $(6, -8)$ lies on the circle.


$(5, 9)$
 $x^2 + y^2 = 100$
 $5^2 + 9^2$
 $= 25 + 81$
 $= 106$
 \therefore Since $106 > 100$
 $(5, 9)$ lies outside the circle.

Circles with Centres *Not* at the Origin

The radius in the above circle is the distance between (x, y) and the origin, i.e. $r = \sqrt{(x-0)^2 + (y-0)^2}$. If we move the centre to the point (a, b) , then $r = \sqrt{(x-a)^2 + (y-b)^2}$.

Squaring both sides, we can now also say that:

The equation $(x - a)^2 + (y - b)^2 = r^2$ describes a circle with centre (a, b) and radius r



Example 3: Write down the centre and radius of each circle.

a) $(x - 1)^2 + (y + 3)^2 = 4$

b) $(x + 9)^2 + (y - 2)^2 = 20$

c) $(x - 5)^2 + y^2 = 400$

Example 4: A is the point (4, 9) and B is the point (-2, 1).
Find the equation of the circle for which AB is the diameter.

$$MP_{AB} = \left(\frac{4+(-2)}{2}, \frac{9+1}{2} \right) = (1, 5)$$

$$radius = 5$$


$$d = \sqrt{(9-1)^2 + (4-(-2))^2}$$

$$d = \sqrt{16+9}$$

$$d = \sqrt{25}$$

$$d = 5$$

$$(x-1)^2 + (y-5)^2 = 25$$



Example 5: Points P, Q and R have coordinates (-10, 2), (5, 7) and (6, 4) respectively.

a) Show that triangle PQR is right angled at Q.

$$\vec{QR} = r - q = \begin{pmatrix} 6 \\ 4 \\ -3 \end{pmatrix} - \begin{pmatrix} 5 \\ 7 \\ 7 \end{pmatrix} = \begin{pmatrix} 1 \\ -3 \\ -3 \end{pmatrix}$$

$$\vec{QP} = p - q = \begin{pmatrix} -10 \\ 2 \\ 7 \end{pmatrix} - \begin{pmatrix} 5 \\ 7 \\ 7 \end{pmatrix} = \begin{pmatrix} -15 \\ -5 \\ 0 \end{pmatrix}$$

$$\vec{QR} \cdot \vec{QP} = 1(-15) + (-3)(-5) + (-3)(0) = -15 + 15 = 0$$

\therefore Since $\vec{QR} \cdot \vec{QP} = 0$
lines are perpendicular
 $\therefore \Delta PQR$ is right angled at Q.

b) Hence find the equation of the circle passing through points P, Q and R.

$$MP_{PR} = (-2, 3)$$

$$d = \sqrt{(3-4)^2 + (-2-6)^2}$$

$$d = \sqrt{1+64}$$

$$d = \sqrt{65} \quad radius = \sqrt{65}$$

$$(x+2)^2 + (y-3)^2 = 65$$

The General Equation of a Circle

For the circle described in Example 3a, we could expand the brackets and simplify to obtain the equation $x^2 + y^2 - 2x + 6y + 6 = 0$, which would also describe a circle with centre (1, -3) and radius 2.

$$\text{For } x^2 + y^2 + 2gx + 2fy + c = 0,$$

$$(x^2 + 2gx) + (y^2 + 2fy) = -c$$

$$(x^2 + 2gx + g^2) + (y^2 + 2fy + f^2) = g^2 + f^2 - c$$

$$(x + g)^2 + (y + f)^2 = (g^2 + f^2 - c)$$

Therefore, the circle described by

$x^2 + y^2 + 2gx + 2fy + c = 0$

has centre $(-g, -f)$ and $r = \sqrt{g^2 + f^2 - c}$
--

Example 6: Find the centre and radius of the circle with equation $x^2 + y^2 - 4x + 8y - 5 = 0$

$$\text{Centre } (2, -4)$$

$$\begin{aligned} \text{radius} &= \sqrt{4 + 16 - (-5)} \\ &= \sqrt{25} \\ &= \underline{\underline{5}} \end{aligned}$$

Example 7: State why the equation $x^2 + y^2 - 4x - 4y + 15 = 0$ does not represent a circle.

$$\text{Centre } (2, 2)$$

$$\begin{aligned} r &= \sqrt{4 + 4 + (-15)} \\ &= \sqrt{-7} \end{aligned}$$

Since we can't $\sqrt{-ve}$.

Example 8: State the range of values of c such that the equation $x^2 + y^2 - 4x + 6y + c = 0$ describes a circle.

Centre $(2, -3)$

$$r = \sqrt{4 + 9 - c}$$


$$r = \sqrt{13 - c}$$

$$13 - c > 0$$

$$-c > -13$$

$$\underline{c < 13}$$

Example 9: Find the equation of the circle concentric with $x^2 + y^2 + 6x - 2y - 54 = 0$ but with radius half its size.



Centre $(-3, 1)$

$$r = \sqrt{9 + 1 - (-54)}$$

$$r = \sqrt{64}$$

$$r = 8$$

$$(-3, 1) \quad r = 4$$




$$(x+3)^2 + (y-1)^2 = 16 \quad \checkmark$$

$$(x-a)^2 + (y-b)^2 = r^2$$

Intersection of Lines and Circles

As with parabolas, there are **three** possibilities when a line and a circle come into contact, and we can examine the roots of a rearranged quadratic equation to determine which has occurred. However:

We CANNOT make the circle and line equations equal to each other: the line equation must be substituted INTO the circle equation to obtain our quadratic equation!

		
<p>Two points of contact 2 distinct roots $b^2 - 4ac > 0$</p>	<p>One point of contact Equal roots $b^2 - 4ac = 0$</p>	<p>No points of contact No real roots $b^2 - 4ac < 0$</p>

As with parabolas, the most common use of this technique is to show tangency.

Example 10: Find the coordinates of the points of intersection of the line $y = 2x - 1$ and the circle $x^2 + y^2 - 2x - 12y + 27 = 0$.

"y = y"

$$x^2 + (2x-1)^2 - 2x - 12(2x-1) + 27 = 0$$

$$x^2 + 4x^2 - 4x + 1 - 2x - 24x + 12 + 27 = 0$$

$$5x^2 - 30x + 40 = 0$$

$$5(x^2 - 6x + 8) = 0 \quad \begin{matrix} 3x-4 \\ x-2 \end{matrix}$$

$$5(x-4)(x-2) = 0$$

"y = y"

When $x = 4$
 $y = 2(4) - 1$
 $y = 7$

When $x = 2$
 $y = 2(2) - 1$
 $y = 3$

$\therefore (4, 7) \text{ \& } (2, 3)$

Example 11: Show that the line $y = 3x + 10$ is a tangent to the circle $x^2 + y^2 - 8x - 4y - 20 = 0$ and establish the coordinates of the point of contact.

"y = y"

$$x^2 + (3x+10)^2 - 8x - 4(3x+10) - 20 = 0$$

$$x^2 + 9x^2 + 60x + 100 - 8x - 12x - 40 - 20 = 0$$

$$10x^2 + 40x + 40 = 0$$

$$10(x^2 + 4x + 4) = 0 \quad \begin{matrix} x+2 \\ x+2 \end{matrix}$$

$$10(x+2)^2 = 0$$

"y = y"

When $x = 2$
 $y = 2(2) - 1$
 $y = 3$


\therefore Since equal roots, line is a tangent to the circle.

$$x = -2$$

When $x = -2$, $y = 3(-2) + 10$

\therefore PoC $(-2, 4)$ $y = 4$

Example 12: Find the equations of the tangents to the circle $x^2 + y^2 = 9$ from the point $(0, 5)$.



"y = y"

$$x^2 + (mx + 5)^2 = 9$$

$$x^2 + m^2x^2 + 10mx + 25 = 9$$

$$x^2 + m^2x^2 + 10mx + 16 = 0.$$

Tangency when $b^2 - 4ac = 0$.

$$x^2(1+m^2) + 10mx + 16 = 0.$$

$$(10m)^2 - 4(1+m^2)16 = 0.$$

$$100m^2 - 64 - 64m^2 = 0$$

$$36m^2 = 64$$

$$m^2 = \frac{64}{36}$$

$$m = \pm \sqrt{\frac{64}{36}}$$

$$m = \pm \frac{8}{6}$$

$$m = \pm \frac{4}{3}$$

$$y = \frac{4}{3}x + 5$$

$$y = -\frac{4}{3}x + 5$$


$a = 1+m^2$
 $b = 10m$
 $c = 16$

Tangents to Circles at Given Points

Remember: at the point of contact, the radius and tangent meet at 90° (i.e., they are perpendicular).

To find a tangent at a given point:

- Find the centre of the circle
- Find the gradient of the radius (joining C and the given point)
- Find the gradient of the tangent (flip and make negative)
- Sub the gradient and the original point into $y - b = m(x - a)$




Example 13: Find the equation of the tangent to $x^2 + y^2 - 14x + 6y - 87 = 0$ at the point $(-2, 5)$.

Past Paper Example 1: A circle has centre $C(-2, 3)$ and passes through point $P(1, 6)$.

a) Find the equation of the circle.

b) PQ is a diameter of the circle. Find the equation of the tangent to this circle at Q .



Past Paper Example 2:

a) Show that the line with equation $y = 3 - x$ is a tangent to the circle with equation

$$x^2 + y^2 + 14x + 4y - 19 = 0$$


and state the coordinates of P, the point of contact.

b) Relative to a suitable set of coordinate axes, the diagram opposite shows the circle from a) and a second smaller circle with centre C.

The line $y = 3 - x$ is a common tangent at the point P.

The radius of the larger circle is three times the radius of the smaller circle.

Find the equation of the smaller circle.



Past Paper Example 3: Given that the equation


$$x^2 + y^2 - 2px - 4py + 3p + 2 = 0$$

represents a circle, determine the range of values of p .

Past Paper Example 4: Circle P has equation $x^2 + y^2 - 8x - 10y + 9 = 0$. Circle Q has centre $(-2, -1)$ and radius $2\sqrt{2}$.

- a) i) Show that the radius of circle P is $4\sqrt{2}$.
ii) Hence show that circles P and Q touch.

- b) Find the equation of the tangent to circle Q at the point $(-4, 1)$




Quadratic Functions

Finding the Equation of a Quadratic Function From Its Graph: $y = k(x - a)(x - b)$

If the graph of a quadratic function has roots at $x = -1$ and $x = 5$, a reasonable guess at its equation would be $y = x^2 - 4x - 5$, i.e. from $y = (x + 1)(x - 5)$.

However, as the diagram shows, there are many parabolas which pass through these points, all of which belong to the family of functions $y = k(x + 1)(x - 5)$.

To find the equation of the original function, we need the roots and one other point on the curve (to allow us to determine the value of k).



Example 1: State the equation of the graph below in the form $y = ax^2 + bx + c$.

$$y = k(x-a)(x-b)$$

$$y = k(x-1)(x-3)$$


When $x=0$ & $y=6$

$$6 = k(0-1)(0-3)$$

$$6 = k(-1)(-3)$$

$$3k = 6$$

$$k = 2$$



$$y = 2(x-1)(x-3)$$

$$y = 2(x^2 - 4x + 3)$$

$$y = 2x^2 - 8x + 6$$

$$y = (2x-2)(x-3)$$

$$y = 2x^2 - 2x - 6x + 6$$

$$y = 2x^2 - 8x + 6.$$


Completing the Square (Revision)

The diagram shows the graphs of two quadratic functions.

If the graph of $y = x^2$ is shifted q units to the right, followed by r units up, then the graph of $y = (x - q)^2 + r$ is obtained.

As the turning point of $y = x^2$ is $(0, 0)$, it follows that the new curve has a turning point at (q, r) .


A quadratic equation written as $y = p(x - q)^2 + r$ is said to be in the **completed square form**.



Example 2: (i) Write the following in the form $y = (x + q)^2 + r$ and find the minimum value of y .
(ii) Hence state the minimum value of y and the corresponding value of x .

a) $y = x^2 + 6x + 10$

b) $y = x^2 - 3x + 1$



Completing the Square when the x^2 Coefficient $\neq 1$

Example 3: Write $y = 3x^2 + 12x + 5$ in the form
 $y = p(x + q)^2 + r$.

Example 4: Write $y = 5 + 12x - x^2$ in the form
 $y = p - (x + q)^2$.

$$\begin{aligned}y &= -x^2 + 12x + 5 \\y &= -(x^2 - 12x) + 5 \\y &= -(x - 6)^2 + 5 + 36 \\y &= -(x - 6)^2 + 41 \\y &= 41 - (x - 6)^2\end{aligned}$$

Example 5:

- a) Write $y = x^2 - 10x + 28$ in the form
 $y = (x + p)^2 + q$.

$$y = (x-5)^2 + 28 - 25$$
$$y = (x-5)^2 + 3$$

- b) Hence find the maximum value of $\frac{18}{x^2 - 10x + 28}$

$$\frac{18}{(x-5)^2 + 3}$$

When $x = 5$

$$\frac{18}{(5-5)^2 + 3}$$
$$= \frac{18}{3}$$

$$= 6.$$

Max value = 6 when $x = 5$

Solving Quadratic Equations via Completing the Square

Quadratic equations which do not easily factorise can be solved in two ways: (i) completing the square, or (ii) using the quadratic formula. In fact, both methods are essentially the same, as the quadratic formula is obtained by solving $y = ax^2 + bx + c$ via completing the square.

Example 6: State the **exact** values of the roots of the equation $2x^2 - 4x + 1 = 0$ by:

a) using the quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad \begin{array}{l} a=2 \\ b=-4 \\ c=1 \end{array}$$

$$x = \frac{-(-4) \pm \sqrt{(-4)^2 - 4 \times 2 \times 1}}{2 \times 2}$$

$$x = \frac{4 \pm \sqrt{8}}{4}$$

$$x = \frac{4 + \sqrt{8}}{4}, \frac{4 - \sqrt{8}}{4}$$

$$x = \underline{\underline{1.7}}, \underline{\underline{0.29}}$$

b) completing the square

$$2(x^2 - 2x) + 1 = 0$$

$$2(x-1)^2 + 1 - 2 = 0$$

$$2(x-1)^2 - 1 = 0$$

$$2(x-1)^2 = 1$$

$$(x-1)^2 = \frac{1}{2}$$

$$x-1 = \pm \sqrt{\frac{1}{2}}$$

$$x = \pm \frac{1}{\sqrt{2}} + 1$$

$$x = \frac{1}{\sqrt{2}} + 1, -\frac{1}{\sqrt{2}} + 1$$

Solving Quadratic Inequations

Quadratic inequations are easily solved by making a sketch of the equivalent quadratic function, and determining the regions above or below the x - axis.


Example 7: Find the values of x for which: a) $2x^2 - 7x + 6 > 0$ b) $2x^2 - 7x + 6 < 0$

First, sketch $y = 2x^2 - 7x + 6$

$$2x^2 - 7x + 6 = 0$$

$$(2x-3)(x-2) = 0$$

$$x = \frac{3}{2}, x = 2$$



a) $2x^2 - 7x + 6 > 0$
 $x < \frac{3}{2} \text{ \& } x > 2$

b) $2x^2 - 7x + 6 < 0$
 $\frac{3}{2} < x < 2$

$x > \frac{3}{2}$
 $x < 2$
 $\frac{3}{2} < x < 2$

Roots of Quadratic Equations and The Discriminant (Revision)

For $y = ax^2 + bx + c$, $b^2 - 4ac$ is known as the **discriminant**.

- $b^2 - 4ac > 0$ gives real, unequal roots
- $b^2 - 4ac = 0$ gives real, equal roots
- $b^2 - 4ac < 0$ gives NO real roots

If $b^2 - 4ac$ gives a perfect square, the roots are RATIONAL
If $b^2 - 4ac$ does NOT give a perfect square, the roots are IRRATIONAL (i.e. surds)

Example 8: Determine the nature of the roots of the equation $4x(x - 3) = 9$




Example 9: Find the value(s) of p given that $2x^2 + 4x + p = 0$ has real roots.

Example 10: Find the value(s) of r given that $x^2 + (r - 3)x + r = 0$ has no real roots.

Tangents to Curves: Using the Discriminant

To find the points of contact between a line and a curve, we make the curve and line equations equal (i.e. make $y = y$) to obtain a quadratic equation, and solve to find the x -coordinates.

By finding the discriminant of this quadratic equation, we can work out **how many** points of contact there are between the line and the curve. There are 3 options:

		
Two points of contact 2 distinct roots $b^2 - 4ac > 0$	One point of contact Equal roots $b^2 - 4ac = 0$	No points of contact No real roots $b^2 - 4ac < 0$

The most common use for this technique is to show that a line is a tangent to a curve

Example 11: Show that the line $y = 3x - 13$ is a tangent to the curve $y = x^2 - 7x + 12$, and find the coordinates of the point of contact.

$$\begin{aligned}
 y &= y \\
 x^2 - 7x + 12 &= 3x - 13 \\
 x^2 - 10x + 25 &= 0 \quad x-5 \\
 (x-5)^2 &= 0 \\
 \therefore \text{Since we have equal roots, line is a tangent.} \\
 x &= 5 \\
 \text{When } x=5, y &= 3(5) - 13 \\
 y &= 2 \\
 \therefore \text{P.C. } &\underline{(5, 2)}
 \end{aligned}$$

Example 12: Find two values of m such that $y = mx - 7$ is a tangent to $y = x^2 + 2x - 3$

$$y = y.$$

$$x^2 + 2x - 3 = mx - 7$$

$$x^2 + 2x - mx - 3 + 7 = 0.$$

$$x^2 + 2x - mx + 4 = 0.$$

Tangency occurs when $b^2 - 4ac = 0$.

$$x^2 + x(2-m) + 4 = 0$$

$$a = 1$$

$$(2-m)^2 - 4 \times 1 \times 4 = 0$$

$$b = 2-m$$

$$c = 4.$$

$$4 - 4m + m^2 - 16 = 0$$

$$m^2 - 4m - 12 = 0 \quad m \quad -6$$


$$(m-6)(m+2) = 0$$

$$\underline{\underline{m=6}}, \underline{\underline{m=-2}}$$

Past Paper Example 1: Express $2x^2 + 12x + 1$ in the form $a(x + b)^2 + c$.

$$\begin{aligned} & 2(x^2 + 6x) + 1 \\ &= 2(x+3)^2 + 1 - 18 \\ &= \underline{2(x+3)^2 - 17} \end{aligned}$$

Past Paper Example 2: Given that $2x^2 + px + p + 6 = 0$ has no real roots, find the range of values for p .



No real roots when $b^2 - 4ac < 0$

$$\begin{aligned} p^2 - 4 \times 2 \times (p+6) &< 0 \\ p^2 - 8p - 48 &< 0 \end{aligned}$$

$$-4 < p < 12$$

$$12 > p > -4$$

$$\begin{aligned} a &= 2 \\ b &= p \\ c &= p+6. \end{aligned}$$

$$\begin{aligned} & p^2 - 8p - 48 = 0 \\ & (p-12)(p+4) = 0 \\ & \underline{p=12}, \underline{p=-4} \\ & p > -4 \\ & -4 < p. \end{aligned}$$

Past Paper Example 3: Show that the roots of $(k-2)x^2 - 3kx + 2k = -2x$ are always real.

Real roots occur when $b^2 - 4ac \geq 0$.

$$(k-2)x^2 - 3kx + 2k = -2x$$

$$(k-2)x^2 + x(-3k+2) + 2k = 0$$

$$(a-3k)^2 - 4(k-2)(2k) \quad \begin{array}{l} a = k-2 \\ b = -3k+2 \\ c = 2k \end{array}$$

$$4 - 12k + 9k^2 - 8k^2 + 16k$$

$$k^2 + 4k + 4 \quad \frac{k^2}{k^2}$$


$$(k+2)^2$$

$$\therefore \text{Since } (k+2)^2 \geq 0 \text{ for all } k, \quad \begin{array}{l} (a-b)^2 \\ = a^2 - 2ab + b^2 \end{array} \quad |$$

roots are always real.


Calculus 2: Integration

The reverse process to differentiation is known as integration.



As it is the opposite of finding the derivative, the function obtained by integration is sometimes called the **anti-derivative**, but is more commonly known as the **integral**, and is given the sign \int .

If $f(x) = x^n$, then $\int x^n dx$ is "the integral of x^n with respect to x "



In general:

$$\int ax^n dx = \frac{ax^{n+1}}{n+1} + C \quad (n \neq -1)$$

- INtegration INcreases the power!**
1. Write as ax^n
 2. Increase the power by 1
 3. Divide by the new power

Example 1: Find (remember "+C"):

$$\text{a) } \int 2x \, dx$$

$$\underline{\underline{x^2 + C}}$$

$$\text{b) } \int 4t^2 \, dt$$

$$= \frac{4t^3}{3} + C$$

$$\text{c) } \int (3x^5 - 4) \, dx$$

$$= \frac{3x^6}{6} - 4x + C$$

$$\text{d) } \int \frac{3}{g^4} \, dg \quad (g \neq 0)$$

$$= \int 3g^{-4} \, dg$$

$$\frac{3g^{-3}}{-3} + C$$

$$= -g^{-3} + C$$

$$= \frac{-1}{g^3} + C$$

$$\text{e) } \int 6\sqrt[5]{p^3} \, dp$$

$$\int 6p^{3/5} \, dp$$

$$= \frac{6p^{8/5}}{8/5} + C$$

$$= \frac{5}{8} \times 6p^{8/5} + C$$

$$= \frac{15}{4} p^{8/5} + C$$

$$= \underline{\underline{\frac{15}{4} \sqrt[5]{p^8} + C}}$$

$$\text{f) } \int \frac{4y-3}{y^{2/3}} \, dy \quad (y \neq 0)$$

$$\int 4y^{1/3} - 3y^{-2/3} \, dy$$

$$= \frac{4y^{4/3}}{4/3} - \frac{3y^{1/3}}{1/3} + C$$

$$= 3y^{4/3} - 9y^{1/3} + C$$

The Definite Integral

A **definite integral** of a function is the difference between the integrals of $f(x)$ at two values of x . Suppose we integrate $f(x)$ and get $F(x)$. Then the integral of $f(x)$ when $x = a$ would be $F(a)$, and the integral when $x = b$ would be $F(b)$.

The definite integral of $f(x)$, with respect to x , between a and b , is written as:

$$\int_a^b f(x)dx = F(b) - F(a) \quad (\text{where } b > a)$$

For example, the integral of $f(x) = 2x^2 - 4$ between the values $x = -3$ and $x = 5$ is written as

$$\int_{-3}^5 (2x^2 - 4)dx \text{ and reads "the integral from -3 to 5 of } 2x^2 - 4 \text{ with respect to } x \text{"}$$

Note: definite integrals do NOT include the constant of integration!

$$\int_a^b f(x) = [F(b) + C] - [F(a) + C] = F(b) - F(a)$$

Example 2: Evaluate $\int_{-1}^3 (2x-1)dx$

$$\begin{aligned} &= \left[\frac{2x^2}{2} - x \right]_{-1}^3 \\ &= [x^2 - x]_{-1}^3 \\ &= [3^2 - 3] - [(-1)^2 - (-1)] \\ &= 6 - 2 = 4 \end{aligned}$$

Always -ve

- To find a definite integral:
- prepare the function for integration
 - integrate as normal, but write inside square brackets with the limits to the right
 - sub each limit into the integral, and subtract the integral with the lower limit from the one with the higher limit

Example 3: Evaluate $\int_0^2 (p+1)(p-1)dp$

$$\begin{aligned} &= \int_0^2 (p^2 - 1) dp \\ &= \left[\frac{p^3}{3} - p \right]_0^2 \\ &= \left[\frac{2^3}{3} - 2 \right] - \left[\frac{0^3}{3} - 0 \right] \\ &= \frac{8}{3} - 2 \\ &= \frac{2}{3} \end{aligned}$$

Example 4: Evaluate $\int_1^{\sqrt{3}} (x^2 - 2x)dx$

$$\begin{aligned} &= \left[\frac{x^3}{3} - x^2 \right]_1^{\sqrt{3}} \\ &= \left[\frac{(\sqrt{3})^3}{3} - (\sqrt{3})^2 \right] - \left[\frac{1^3}{3} - 1^2 \right] \\ &= \frac{3\sqrt{3}}{3} - 3 - \left[\frac{1}{3} - 1 \right] \\ &= \frac{\cancel{3}\sqrt{3}}{\cancel{3}} - 3 + \frac{2}{3} \\ &= \sqrt{3} - 3 + \frac{2}{3} \\ &= \sqrt{3} - \frac{7}{3} \end{aligned}$$

Example 5: Find the value of g such that $\int_{-2}^g (6x+5) dx = 6$.

$$\left[\frac{6x^2}{2} + 5x \right]_{-2}^g = 6.$$

$$\left[3x^2 + 5x \right]_{-2}^g = 6$$

$$\left[3g^2 + 5g \right] - \left[3(-2)^2 + 5(-2) \right] = 6$$

$$3g^2 + 5g - [12 - 10] = 6$$

$$3g^2 + 5g - 8 = 0. \quad \begin{matrix} 3g^2 + 8 \\ g - 1 \end{matrix}$$


$$(3g+8)(g-1) = 0$$

$$g = \underline{\underline{-\frac{8}{3}}}, \quad g = \underline{\underline{1}}$$

$$\underline{\underline{\text{Discard}}} \quad \therefore g = \underline{\underline{1}}$$

Ex 9L
Q4

Area Between a Curve and the x - axis.



In the diagram opposite, the area of the shaded section can be obtained by finding the area under the graph from 0 to b, and subtracting the area from 0 to a.



The value of each of these areas can be determined by integrating the function and substituting b or a respectively.

The area enclosed by the curve $y = f(x)$, the lines $y = a$, $y = b$ and the x - axis is equal to the definite integral of $f(x)$ between a and b

$$\text{i.e. Area} = \int_a^b f(x) dx$$

Example 6: For each graph below,

- (i) write down the integrals which describe the shaded regions
- (ii) calculate the area of the shaded region




NOTE: Example 6b shows that areas UNDER the x - axis give NEGATIVE values!

Example 7:

a) Evaluate $\int_{-1}^7 (2x-6) dx$

b) (i) Sketch below the area described by the integral $\int_{-1}^7 (2x-6) dx$.




The answers for 5a and 5b do not match! This is because the area below the axis and the area above cancel each other out (as in 4b, areas below the x - axis give negative values).


To find the area between a curve and the x-axis:

1. Determine the limits which describe the sections above and below the axis
2. Calculate areas separately
3. Find the total, *IGNORING THE NEGATIVE VALUE OF THE SECTION BELOW THE AXIS.*


Example 8: Determine the area of the regions bounded by the curve $y = x^2 - 4x + 3$ and the x - and y - axes.




Example 9: Write down the integrals used to determine the areas shown below:



$$\begin{aligned}
 y &= y \\
 6x - x^2 &= 5 \\
 x^2 - 6x + 5 &= 0 \quad x-5 \\
 (x-5)(x-1) &= 0 \\
 x &= 5, x=1 \\
 \int_1^5 (6x - x^2 - 5) dx.
 \end{aligned}$$




$$\begin{aligned}
 2x^2 - 3 &= x \\
 2x^2 - x - 3 &= 0 \quad 2x-3 \\
 (2x-3)(x+1) &= 0 \\
 x &= \frac{3}{2}, x=-1 \\
 \int_{-1}^{3/2} (x - (2x^2 - 3)) dx \\
 \int_{-1}^{3/2} (x - 2x^2 + 3) dx.
 \end{aligned}$$



$$\int_0^1 (\sqrt{x} - x^2) dx.$$

Example 10: Find the area enclosed between the curve $y = x^3 - x^2 - 5x$ and the line $y = x$



$$\begin{aligned}
 y &= y \\
 x^3 - x^2 - 5x &= x \\
 x^3 - x^2 - 6x &= 0 \\
 x(x^2 - x - 6) &= 0 \quad x = 0 \\
 x(x-3)(x+2) &= 0 \\
 x &= 0, x=3, x=-2
 \end{aligned}$$

$$\begin{aligned}
 \textcircled{A} \int_{-2}^0 x^3 - x^2 - 5x - x \, dx \\
 &= \int_{-2}^0 x^3 - x^2 - 6x \, dx \\
 &= \left[\frac{x^4}{4} - \frac{x^3}{3} - \frac{6x^2}{2} \right]_{-2}^0 \\
 &= [0] - \left[\frac{-2^4}{4} - \frac{(-2)^3}{3} - \frac{6(-2)^2}{2} \right] \\
 &= 0 - \left[4 + \frac{8}{3} - 12 \right] \\
 &= 5\frac{1}{3} \\
 \therefore A_{\text{A}} &= 5\frac{1}{3} \text{ units}^2
 \end{aligned}$$

$$\begin{aligned}
 \textcircled{B} \int_0^3 x - (x^3 - x^2 - 5x) \, dx \\
 &= \int_0^3 x - x^3 + x^2 + 5x \, dx \\
 &= \int_0^3 6x - x^3 + x^2 \, dx \\
 &= \left[\frac{6x^2}{2} - \frac{x^4}{4} + \frac{x^3}{3} \right]_0^3 \\
 &= \left[\frac{6 \times 3^2}{2} - \frac{3^4}{4} + \frac{3^3}{3} \right] - [0] \\
 &= 27 - \frac{81}{4} + 9 - 0 \\
 &= 15\frac{3}{4} \\
 A_{\text{B}} &= 15\frac{3}{4} \text{ units}^2
 \end{aligned}$$

$$\begin{aligned}
 A_{\text{T}} &= 5\frac{1}{3} + 15\frac{3}{4} \\
 &= 21\frac{1}{2} \text{ units}^2
 \end{aligned}$$

Differential Equations

If we know the derivative of a function (e.g. $f'(x) = 6x^2 - 3$), we can obtain a formula for the original function by integration. This is called a **differential equation**, and gives us the function in terms of x and C (which we can then evaluate if we have a point on the graph of the function).

Example 11: The gradient of a tangent to the curve of $y = f(x)$ is $24x^2 + 10x$, Express y in terms of x , given that the graph of $y = f(x)$ passes through the point $(-1, -10)$.

$$\frac{dy}{dx} = 24x^2 + 10x$$

$$y = \frac{24x^3}{3} + \frac{10x^2}{2} + C$$

$$y = 8x^3 + 5x^2 + C$$

$$-10 = 8(-1)^3 + 5(-1)^2 + C$$

$$-10 = -8 + 5 + C$$

$$C = -7$$


$$\therefore y = 8x^3 + 5x^2 - 7$$

Ex 9Q
Qs 3, 4, 5

Past Paper Example 1: Evaluate $\int_1^9 \frac{x+1}{\sqrt{x}} dx$

$$\begin{aligned}
 & \int_1^9 \frac{x+1}{x^{1/2}} dx \\
 &= \int_1^9 \frac{x^1}{x^{1/2}} + \frac{1}{x^{1/2}} dx \\
 &= \int_1^9 x^{1/2} + x^{-1/2} dx \\
 &= \left[\frac{x^{3/2}}{3/2} + \frac{x^{1/2}}{1/2} \right]_1^9 \\
 &= \left[\frac{2x^{3/2}}{3} + 2x^{1/2} \right]_1^9 \\
 &= \left[\frac{2\sqrt{x^3}}{3} + 2\sqrt{x} \right]_1^9 \\
 &= \left[\frac{2 \times \sqrt{9^3}}{3} + 2\sqrt{9} \right] - \left[\frac{2\sqrt{1^3}}{3} + 2\sqrt{1} \right] \\
 &= [18 + 6] - \left[\frac{2}{3} + 2 \right] \quad 18 + 6 - \frac{2}{3} - 2 \\
 &\quad 22 - \frac{2}{3} \\
 &\quad \underline{21\frac{1}{3}}
 \end{aligned}$$

Past Paper Example 2: Find area enclosed between the curves $y = 1 + 10x - 2x^2$ and $y = 1 + 5x - x^2$.



$$\begin{aligned}
 & y = y \\
 & 1 + 10x - 2x^2 = 1 + 5x - x^2 \\
 & x^2 - 5x = 0 \\
 & x(x - 5) = 0 \\
 & x = 0, x = 5.
 \end{aligned}$$

$$\begin{aligned}
 & \int_0^5 (1 + 10x - 2x^2 - (1 + 5x - x^2)) dx \\
 & = \int_0^5 (1 + 10x - 2x^2 - 1 - 5x + x^2) dx \\
 & = \int_0^5 (5x - x^2) dx \\
 & = \left[\frac{5x^2}{2} - \frac{x^3}{3} \right]_0^5 \\
 & = \left[\frac{5 \times 5^2}{2} - \frac{5^3}{3} \right] - [0] \\
 & = \frac{375}{2} - \frac{125}{3} \\
 & = \frac{125}{6} \text{ units}^2
 \end{aligned}$$


$$\begin{aligned}
 & 375 - 250 \\
 & \frac{125}{6}
 \end{aligned}$$

Past Paper Example 3: The parabola shown in the diagram has equation

$$y = 32 - 2x^2.$$

The shaded area lies between the lines $y = 14$ and $y = 24$

Calculate the shaded area.



$$y = 14$$

$$32 - 2x^2 = 14$$

$$2x^2 = 18$$

$$x^2 = 9$$

$$x = \pm 3.$$

$$y = 24$$

$$32 - 2x^2 = 24$$

$$2x^2 = 8$$

$$x^2 = 4$$

$$x = \pm 2.$$



$$\int_{-3}^3 24 dx.$$

$$\int_{-3}^3 32 - 2x^2 - 14 dx$$

$$= \left[32x - \frac{2x^3}{3} - 14x \right]_{-3}^3$$

$$= \left[18x - \frac{2x^3}{3} \right]_{-3}^3$$

$$= \left[18(3) - \frac{2(3)^3}{3} \right] - \left[18(-3) - \frac{2(-3)^3}{3} \right]$$

$$= 72 \text{ units}^2$$

$$\int_{-2}^2 32 - 2x^2 - 24 dx.$$

$$= \int_{-2}^2 8 - 2x^2 dx$$

$$= \left[8x - \frac{2x^3}{3} \right]_{-2}^2$$

$$= \left[8(2) - \frac{2(2)^3}{3} \right] - \left[8(-2) - \frac{2(-2)^3}{3} \right]$$

$$= \left[16 - \frac{16}{3} \right] - \left[-16 + \frac{16}{3} \right]$$

$$= 16 - \frac{16}{3} + 16 - \frac{16}{3}.$$

$$= 32 - \frac{32}{3}$$

$$= \frac{64}{3} \text{ units}^2.$$

$$A_T = 72 - \frac{64}{3}$$

$$= \frac{152}{3} \text{ units}^2$$

Polynomials

A **polynomial** is an expression with terms of the form ax^n , where n is a whole number.

For example, $5p^4 - 3p^3$ is a polynomial, but $3p^{-1}$ or $\sqrt[3]{p^2}$ are not.

The **degree** of a polynomial is its highest power, e.g. the polynomial above has a degree of 4.

The number part of each term is called its **coefficient**, e.g. the coefficients of p^4 , p^3 and p above are 5, -3 and 0 (as there is no p term!) respectively (note that $5p^4$ would also be a polynomial on its own, with coefficients of zero for all other powers of p).

Evaluating Polynomials

An easy way to find out the value of a polynomial function is by using a **nested table**.

Example 1: Evaluate $f(4)$ for $f(x) = 2x^4 - 3x^3 - 10x^2 - 5x + 7$.

4	2	-3	-10	-5	7	
	↓	↗	↖	↖	↖	
	2	8	20	40	140	
		5	10	35	147	

← Line up coefficients

Example 2: Evaluate $f(-1)$ for $f(x) = 3x^5 - 2x^3 + 4$.

-1	3	0	-2	0	0	4
	↓	↗	↖	↖	↖	
	3	-3	3	-1	1	-1
		-3	1	-1	1	3

Missing powers have coefficients of zero!

Synthetic Division

Dividing 67 by 9 gives an answer of “7 remainder 4”. We can write this in two ways:

$$67 \div 9 = 7 \text{ remainder } 4 \qquad \text{OR} \qquad 9 \times 7 + 4 = 67$$

For this problem, 9 is the **divisor**, 7 is the **quotient**, and 4 is the **remainder** (note that if we were dividing 63 by 9, the remainder would be zero, since 9 is a **factor** of 63).

Example 3:

a) Remove brackets and simplify:

$$\begin{aligned} & (x-3)(x^2+9x-12)+11 \\ & x(x^2+9x-12)-3(x^2+9x-12)+11 \\ & x^3+9x^2-12x-3x^2-27x+36+11 \\ & x^3+6x^2-39x+47 \end{aligned}$$

b) Evaluate f(3) for $f(x) = x^3 + 6x^2 - 39x + 47$

$$\begin{array}{r|rrrr} 3 & 1 & 6 & -39 & 47 \\ & & 3 & 27 & -36 \\ \hline & 1 & 9 & -12 & 11 \end{array}$$

↓
remainder.

This shows that:

$$x^3 + 6x^2 - 39x + 47 = (x-3)(x^2 + 9x - 12) + 11$$

OR

$$\underline{(x^3 + 6x^2 - 39x + 47) \div (x - 3) = (x^2 + 9x - 12) \text{ remainder } 11}$$

Example 4: Find the remainder on dividing $x^3 - x^2 - x + 5$ by $(x + 5)$.

$$\begin{array}{r|rrrr} -5 & 1 & -1 & -1 & 5 \\ & \downarrow & -5 & 30 & -145 \\ \hline & 1 & -6 & 29 & -140 \end{array}$$

Quotient. = $x^2 - 6x + 29$
Remainder = -140 .

$$\begin{aligned} & (x^2 - 6x + 29)(x + 5) - 140 \\ & = x^3 - x^2 - x + 5 \end{aligned}$$

Example 5: Write $4p^4 + 2p^3 - 6p^2 + 3 \div (2p - 1)$ in the form $(ap - b)Q(p) + R$

$$\begin{array}{r|rrrrr} \frac{1}{2} & 4 & 2 & -6 & 0 & 3 \\ & \downarrow & & & & \\ \hline & & 2 & 2 & -2 & -1 \\ & 4 & 4 & -4 & -2 & 2 \end{array}$$

$$(p - \frac{1}{2})(4p^3 + 4p^2 - 4p - 2) + 2$$

$$\div 2$$

$$(2p - 1)(2p^3 + 2p^2 - 2p - 1) + 2$$

Remainder Theorem and Factor Theorem

Considered together, these two theorems allow us to factorise algebraic functions (remember that a factor is a number or term which divides exactly into another, leaving no remainder).

If polynomial $f(x)$ is divided by $(x - h)$, then the remainder is $f(h)$

On division of polynomial $f(x)$ by $(x - h)$, if $f(h) = 0$, then $(x - h)$ is a factor of $f(x)$

In other words, if the result of synthetic division on a polynomial by h is zero, then h is a root of the polynomial, and $(x - h)$ is a factor of it.

Example 6: $f(x) = 2x^3 - 9x^2 + x + 12$.

a) Show that $(x - 4)$ is a factor of $f(x)$.

$$\begin{array}{r|rrrr} 4 & 2 & -9 & 1 & 12 \\ & \downarrow & 8 & -4 & -12 \\ \hline & 2 & -1 & -3 & 0 \end{array}$$

\therefore Since $r=0$, $(x-4)$ is a factor.

b) Hence factorise $f(x)$ fully.

$$\begin{aligned} & (x-4)(2x^2 - x - 3) \quad \begin{matrix} 2x-3 \\ x+1 \end{matrix} \\ & (x-4)(2x-3)(x+1) \end{aligned}$$

Example 7: Factorise fully $3x^3 + 2x^2 - 12x - 8$.

$$\begin{array}{r|rrrr} 1 & 3 & 2 & -12 & -8 \\ & & 3 & 5 & -7 \\ \hline & 3 & 5 & -7 & -15 \end{array}$$

$$\begin{array}{r|rrrr} 4 & 3 & 2 & -12 & -8 \\ & & 12 & 50 & 152 \\ \hline & 3 & 14 & 44 & 144 \end{array}$$

$$\begin{array}{r|rrrr} -2 & 3 & 2 & -12 & -8 \\ & & -6 & 8 & 8 \\ \hline & 3 & -4 & -4 & 0 \end{array}$$

$$\begin{aligned} & (x+2)(3x^2 - 4x - 4) \quad \begin{matrix} 3x-2 \\ x+2 \end{matrix} \\ & (x+2)(3x+2)(x-2) \end{aligned}$$

Example 8: Find the value of k for which $(x + 3)$ is a factor of $x^3 - 3x^2 + kx + 6$

$$\begin{array}{r|rrrr}
 -3 & 1 & -3 & k & 6 \\
 & \downarrow & -3 & 18 & -3k-54 \\
 \hline
 & 1 & -6 & 18+k & \underline{-3k-48}
 \end{array}$$

Since $(x+3)$ is a factor

$$-3k - 48 = 0.$$

$$-3k = 48$$

$$k = -16$$

$$\underline{\underline{k = -16}}$$

Example 9: Find the values of a and b if $(x - 3)$ and $(x + 5)$ are both factors of $x^3 + ax^2 + bx - 15$

$$\begin{array}{r|rrrr} 3 & 1 & a & b & -15 \\ & & 3 & 9+3a & 27+9a+3b \\ \hline & 1 & 3+a & 9+3a+b & 12+9a+3b \end{array}$$

Since $(x-3)$ is a factor $12 + 9a + 3b = 0$.

$$\begin{array}{r|rrrr} -5 & 1 & a & b & -15 \\ & & -5 & 25-5a & 25a-5b-125 \\ \hline & 1 & a-5 & 25-5a+b & 25a-5b-140 \end{array}$$

Since $(x+5)$ is a factor $25a - 5b - 140 = 0$

$$\begin{array}{r} 9a + 3b = -12 \quad \times 5 \\ 25a - 5b = 140 \quad \times 3 \end{array}$$

$$\begin{array}{r} 45a + 15b = -60 \quad \text{Add!} \\ 75a - 15b = 420 \quad \underline{\underline{\quad}} \\ \hline 120a = 360 \end{array}$$

$$\begin{array}{r} 120a = 360 \\ a = 3 \end{array}$$

$$\begin{array}{r} \text{When } a=3, \quad 45(3) + 15b = -60 \\ 135 + 15b = -60 \end{array}$$

$$15b = -195$$


$$b = -13$$

$$\therefore \underline{\underline{a=3}}, \underline{\underline{b=-13}}$$

Solving Polynomial Equations

Polynomial equations are solved in exactly the same way as we solve quadratic equations: make the right hand side equal to zero, factorise, and solve to find the roots.

Example 10: The graph of the function $y = x^3 - 7x^2 + 7x + 15$ is shown.
Find the coordinates of points A, B and C.



$$\begin{array}{l} 1, 15 \\ 3, 5 \\ -1, -15 \\ -3, -5 \end{array}$$

Curves cut x -axis when $y = 0$

$$x^3 - 7x^2 + 7x + 15 = 0$$

$$\begin{array}{r|rrrr} 3 & 1 & -7 & 7 & 15 \\ & \downarrow & & & \\ & 3 & -12 & -15 & \\ \hline & 1 & -4 & -5 & 0 \end{array}$$

$$(x-3)(x^2 - 4x - 5) = 0$$

$$(x-3)(x-5)(x+1) = 0$$

$$x=3, x=5, x=-1$$


8

Finding a Function from its Graph

This uses exactly the same system as that for quadratic graphs, but with more brackets (see page 19).

Remember: tangents to the x - axis have repeated roots!

Example 11: Find an expression for cubic function $f(x)$.



$$y = k(x-a)(x-b)(x-c)$$

$$y = k(x+2)(x-5)(x-5)$$

When $x=0$ & $y=5$

$$5 = k(0+2)(0-5)(0-5)$$

$$5 = 50k$$

$$k = \frac{1}{10}$$

$$\therefore y = \frac{1}{10}(x+2)(x-5)^2$$

Sketching Polynomial Functions

Example 12: a) Find the x - and y - intercepts of the graph of $y = x^4 - 6x^3 + 13x^2 - 12x + 4$.

Curve cuts x-axis when $y=0$. $x^4 - 6x^3 + 13x^2 - 12x + 4 = 0$.

$$(x-2)(x^3 - 4x^2 + 5x - 2) = 0$$

$$(x-2)(x-1)(x^2 - 3x + 2) = 0 \quad \begin{matrix} x-2 \\ x-1 \end{matrix}$$

$$(x-2)(x-1)(x-2)(x-1) = 0$$

$x=1, x=2$ (1,0) (2,0)
(0,4)

$$\begin{array}{r|rrrrr} 2 & 1 & -6 & 13 & -12 & 4 \\ & & 2 & -8 & 10 & -4 \\ \hline & 1 & -4 & 5 & -2 & 0 \end{array}$$

$$\begin{array}{r|rrrr} 1 & 1 & -4 & 5 & -2 \\ & & 1 & -3 & 2 \\ \hline & 1 & -3 & 2 & 0 \end{array}$$

b) Find the position and nature of the stationary points of $y = x^4 - 6x^3 + 13x^2 - 12x + 4$.

$$\frac{dy}{dx} = 4x^3 - 18x^2 + 26x - 12$$

SP's when $\frac{dy}{dx} = 0$

$$4x^3 - 18x^2 + 26x - 12 = 0$$

$$\begin{array}{r|rrrrr} 1 & 4 & -18 & 26 & -12 \\ & & 4 & -14 & 12 \\ \hline & 4 & -14 & 12 & 0 \end{array}$$

$$(x-1)(4x^2 - 14x + 12) = 0$$

$$2(x-1)(2x^2 - 7x + 6) = 0 \quad \begin{matrix} 2x-3 \\ x-2 \end{matrix}$$


$$2(x-1)(2x-3)(x-2) = 0$$

$x=1, x=\frac{3}{2}, x=2$

x	$\rightarrow 0$	$\rightarrow 1$	$\rightarrow \frac{3}{2}$	$\rightarrow 2$	$\rightarrow 3$
$\frac{dy}{dx}$	$-$	0	$+$	0	$+$
Shape	$ $	$-$	$ $	$-$	$ $

MinTP @ (1, 0)
4 (2, 0)
MaxTP @ ($\frac{3}{2}, \frac{1}{16}$)


c) Hence, sketch and annotate the graph of $y = x^4 - 6x^3 + 13x^2 - 12x + 4$.




Trigonometry: Addition Formulae and Equations

What you must know from National 5!!!

Sine Graph



Cosine Graph



We can use the above graphs to find the values of:

$$\sin 0^\circ = 0$$

$$\sin 90^\circ = 1$$

$$\sin 180^\circ = 0$$

$$\sin 270^\circ = -1$$

$$\sin 360^\circ = 0$$

$$\cos 0^\circ = 1$$


$$\cos 90^\circ = 0$$

$$\cos 180^\circ = -1$$

$$\cos 270^\circ = 0$$

$$\cos 360^\circ = 1$$


We can use these graphs to solve the following:		
$\sin x^\circ = 0$ $(0 \leq x \leq 360)$ $x = 0^\circ, 180^\circ, 360^\circ$	$\sin x^\circ = -1$ $(0 \leq x \leq 360)$ $x = 270^\circ$	$\sin x^\circ = 1$ $(0 \leq x \leq 360)$ $x = 90^\circ$
$\cos x^\circ = 0$ $(0 \leq x \leq 360)$ $x = 90^\circ, 270^\circ$	$\cos x^\circ = -1$ $(0 \leq x \leq 360)$ $x = 180^\circ$	$\cos x^\circ = 1$ $(0 \leq x \leq 360)$ $x = 0^\circ, 360^\circ$



Remember , this means that:

- sin 160° would be +
- cos 200° would be -
- tan 200° would be +
- sin 320° would be -
- and so on...

Related Angles



This diagram can be used to find families of related angles.

For example, for $x = 30^\circ$.
The family of related angles would be:
 $30^\circ, 150^\circ, 210^\circ, 330^\circ$

These angles are related since:

$$\begin{aligned}\sin 30^\circ &= 0.5 \\ \sin 150^\circ &= 0.5 \\ \sin 210^\circ &= -0.5 \\ \sin 330^\circ &= -0.5\end{aligned}$$


Note: The sine of these angles have the same numerical value.

Equations	
<p>Example A: $\sin x^\circ = 0.423 \quad (0 \leq x \leq 360)$ $x = \sin^{-1}(0.423)$ $x = 25^\circ \text{ (R.A)}$</p> <p>$x = (0 + 25)^\circ, (180 - 25)^\circ$ $x = 25^\circ, 155^\circ$</p>	<p>Step 1: Consider 0.423</p> <p>Step 2: We know that we can find the other 3 angles in the family $155^\circ, 205^\circ, 335^\circ$</p> <p>Step 3: We only want the angles which will give +ve answers for sin.</p>
<p>Example B: $\cos x^\circ = -0.584 \quad (0 \leq x \leq 360)$ $x = \cos^{-1}(0.584)$ $x = 54.3^\circ \text{ (R.A)}$</p> <p>$x = (180 - 54.3)^\circ, (180 + 54.3)^\circ$ $x = 125.7^\circ, 234.3^\circ$</p>	<p>Step 1: Consider 0.584 (ignore -ve)</p> <p>Step 2: We know that we can find the other 3 angles in the family $125.7^\circ, 234.3^\circ, 305.7^\circ$</p> <p>Step 3: We only want the angles which will give -ve answers for cos.</p>

Radians

If we draw a circle and make a sector with an arc of exactly one radius long, then the angle at the centre of the sector is called a **radian**.

Remember that Circumference = $\pi D = 2\pi r$. This means that there are 2π radians in a full circle.



$360^\circ = 2\pi$ radians
$180^\circ = \pi$ radians

Example 1: Convert:

a) 90° to radians

b) 60° to radians

c) 225° to radians


d) $\frac{\pi}{4}$ radians to degrees

e) $\frac{4\pi}{3}$ radians to degrees


f) $\frac{11\pi}{6}$ radians to degrees

Exact Values

Consider the following triangles:




A right-angled triangle made by halving an equilateral triangle of side 2 units



A right-angled triangle made by halving a square of side 1 unit

Once we have found the lengths of the missing sides (by Pythagoras' Theorem), the following table of values can be constructed:

	0°	30°	45°	60°	90°
	0	$\left(\frac{\pi}{6}\right)$	$\left(\frac{\pi}{4}\right)$	$\left(\frac{\pi}{3}\right)$	$\left(\frac{\pi}{2}\right)$
Sin					
Cos					
Tan					



Example 2: State the exact values of:

a) $\sin 150^\circ$

b) $\tan 315^\circ$



c) $\cos \frac{7\pi}{6}$

d) $\sin (-30)^\circ$

e) $\cos (-120)^\circ$

f) $-\tan \frac{\pi}{4}$

Example 3: The graphs below is of the form $y = a\sin bx + c$ and $y = a\cos bx + c$ respectively. Identify the values of a , b and c in each graph



Addition Formulae

Finding the value of a compound angle is not quite as simple as adding together the values of the component angles, e.g. $\sin 90^\circ \neq \sin 60^\circ + \sin 30^\circ$. The following formulae must be used:

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$


Example 4: Expand each of the following:

a) $\sin(X + Y)$

b) $\sin(Q + 3P)$

Example 5: Find the exact value of $\sin 75^\circ$.

Example 6: A and B are acute angles where $\tan A = \frac{12}{5}$ and $\tan B = \frac{3}{4}$. Find the value of $\sin(A + B)$.



$$\begin{aligned} \cos 120^\circ &= \sin 420^\circ \\ &= -\cos 60^\circ &= \sin 60^\circ \\ &= -\frac{1}{2} &= \frac{\sqrt{3}}{2} \end{aligned}$$

Example 7: Expand each of the following:


a) $\sin(\alpha - \beta)$
 $= \sin \alpha \cos \beta - \cos \alpha \sin \beta$

b) $\sin\left(2B - \frac{2\pi}{3}\right)$


$$\begin{aligned} &= \sin 2B \cos \frac{2\pi}{3} - \cos 2B \sin \frac{2\pi}{3} \\ &= \sin 2B \times \left(-\frac{1}{2}\right) - \cos 2B \times \frac{\sqrt{3}}{2} \\ &= -\frac{\sin 2B}{2} - \frac{\sqrt{3} \cos 2B}{2} \end{aligned}$$

Example 8: In the diagram opposite:
 AC = CD = 2 units, and AB = BC = 1 unit.

Show that $\sin X$ is exactly $\frac{1}{\sqrt{10}}$



$$\begin{aligned} & \sin(A - B) \\ & \sin A \cos B - \cos A \sin B \\ & = \frac{2}{\sqrt{8}} \times \frac{2}{\sqrt{5}} - \frac{2}{\sqrt{8}} \times \frac{1}{\sqrt{5}} \\ & = \frac{4}{\sqrt{40}} - \frac{2}{\sqrt{40}} \\ & = \frac{2}{\sqrt{40}} = \frac{2}{2\sqrt{10}} \\ & = \frac{1}{\sqrt{10}} \end{aligned}$$



$$\begin{aligned} \sqrt{40} &= \sqrt{4 \times 10} \\ &= 2\sqrt{10} \\ A^2 &= B^2 + C^2 \\ A^2 &= 1^2 + 2^2 \end{aligned}$$

$\cos(A + B)$ and $\cos(A - B)$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

Example 9: Expand the following:

a) $\cos(X - Y)$

$$= \cos X \cos Y + \sin X \sin Y.$$

b) $\cos(X + 315^\circ)$

$$\begin{aligned} &= \cos X \cos 315^\circ - \sin X \sin 315^\circ \\ &= \cos X \times \frac{1}{\sqrt{2}} - \sin X \times \left(-\frac{1}{\sqrt{2}}\right) \\ &= \frac{\cos X}{\sqrt{2}} + \frac{\sin X}{\sqrt{2}} \end{aligned}$$

Example 10:

a) Show that $\frac{\pi}{3} - \frac{\pi}{4} = \frac{\pi}{12}$

$$\begin{aligned} &\text{LHS} \\ &\frac{\pi}{3} - \frac{\pi}{4} \\ &\frac{4\pi}{12} - \frac{3\pi}{12} \\ &= \frac{\pi}{12} \quad \text{As Required} \end{aligned}$$

To summarise:

b) Hence find the exact value of $\cos \frac{\pi}{12}$

$$\begin{aligned} &\cos \frac{\pi}{12} \\ &= \cos \left(\frac{\pi}{3} - \frac{\pi}{4} \right) \\ &= \cos \frac{\pi}{3} \cos \frac{\pi}{4} + \sin \frac{\pi}{3} \sin \frac{\pi}{4} \\ &= \frac{1}{2} \times \frac{1}{\sqrt{2}} + \frac{\sqrt{3}}{2} \times \frac{1}{\sqrt{2}} \\ &= \frac{1}{2\sqrt{2}} + \frac{\sqrt{3}}{2\sqrt{2}} \\ &= \frac{\sqrt{3}+1}{2\sqrt{2}} \end{aligned}$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

CXC Q1, 2a, 3, 5a, 6-10
CXC 110 Q1, 2b, 4, 6, 7, 9.

Trigonometric Identities

NOTE: these are important formulae which are not provided in the exam paper formula sheets!

$$\frac{\sin x^\circ}{\cos x^\circ} = \tan x^\circ$$

$$\sin^2 x^\circ + \cos^2 x^\circ = 1$$

Note that due to the second formula, we can also say that:

$$\cos^2 x^\circ = 1 - \sin^2 x^\circ$$

AND

$$\sin^2 x^\circ = 1 - \cos^2 x^\circ$$

To prove that an identity is true, we need to show that the expression on the left hand side of the equals sign can be changed into the expression on the right hand side.

Example 11: Prove that:

a) $\cos^4 \alpha - \sin^4 \alpha = \cos^2 \alpha - \sin^2 \alpha$

$$\begin{aligned} & \text{LHS} \\ & \cos^4 \alpha - \sin^4 \alpha \\ & (\cos^2 \alpha - \sin^2 \alpha)(\cos^2 \alpha + \sin^2 \alpha) \\ & (\cos^2 \alpha - \sin^2 \alpha) \times 1 \\ & = \cos^2 \alpha - \sin^2 \alpha \end{aligned}$$

--

b) $\tan 3\theta + \tan \theta = \frac{\sin 4\theta}{\cos \theta \cos 3\theta}$

$$\begin{aligned} & \text{LHS} \\ & \tan 3\theta + \tan \theta \\ & = \frac{\sin 3\theta}{\cos 3\theta} \times \frac{\sin \theta}{\cos \theta} \\ & = \frac{\sin 3\theta \cos \theta}{\cos \theta \cos 3\theta} + \frac{\sin \theta \cos 3\theta}{\cos \theta \cos 3\theta} \\ & = \frac{\sin 3\theta \cos \theta + \cos 3\theta \sin \theta}{\cos \theta \cos 3\theta} \\ & = \frac{\sin(3\theta + \theta)}{\cos \theta \cos 3\theta} \\ & = \frac{\sin 4\theta}{\cos \theta \cos 3\theta} \end{aligned}$$

$$3c) \quad \cos(360-x) = \cos x$$

$$\begin{aligned} & \text{LHS} \\ & \cos 360^\circ \cos x + \sin 360^\circ \sin x \\ & 1 \times \cos x + 0 \\ & = \underline{\underline{\cos x}} \end{aligned}$$


$$5a) \quad \frac{\sin(\alpha - \beta)}{\cos \alpha \cos \beta} = \tan \alpha - \tan \beta$$

$$\begin{aligned} & \text{LHS} \\ & \frac{\sin \alpha \cos \beta - \cos \alpha \sin \beta}{\cos \alpha \cos \beta} \end{aligned}$$

$$= \frac{\sin \alpha \cancel{\cos \beta}}{\cos \alpha \cancel{\cos \beta}} - \frac{\cancel{\cos \alpha} \sin \beta}{\cancel{\cos \alpha} \cos \beta}$$

$$= \frac{\sin \alpha}{\cos \alpha} - \frac{\sin \beta}{\cos \beta}$$

$$= \tan \alpha - \tan \beta$$



$$\begin{aligned} 2b) \quad \sin(\alpha + \beta) - \sin(\alpha - \beta) &= 2 \cos \alpha \sin \beta \\ &= \sin \alpha \cos \beta + \cancel{\cos \alpha \sin \beta} - (\sin \alpha \cos \beta - \cancel{\cos \alpha \sin \beta}) \\ &= \cancel{\sin \alpha \cos \beta} + \cos \alpha \sin \beta - \cancel{\sin \alpha \cos \beta} + \cos \alpha \sin \beta \\ &= 2 \cos \alpha \sin \beta \end{aligned}$$

$$\begin{aligned} 3d) \quad \cos(180+x) &= -\cos x \\ &= \text{LHS} \\ & \cos 180 \cos x - \sin 180 \sin x \\ &= -1 \times \cos x - 0 \times \sin x \\ &= -\cos x \end{aligned}$$

Double Angle Formulae

$$\begin{aligned}\sin 2A &= \sin(A + A) \\ &= \sin A \cos A + \cos A \sin A \\ &= 2\sin A \cos A\end{aligned}$$

$$\begin{aligned}\cos 2A &= \cos(A + A) \\ &= \cos A \cos A - \sin A \sin A \\ &= \cos^2 A - \sin^2 A.\end{aligned}$$

Since $\cos^2 x^\circ = 1 - \sin^2 x^\circ$ and $\sin^2 x^\circ = 1 - \cos^2 x^\circ$ we can further expand the formula for $\cos 2A$:

$$\begin{aligned}\cos 2A &= \cos^2 A - \sin^2 A \\ &= 1 - \sin^2 A - \sin^2 A \\ &= 1 - 2\sin^2 A\end{aligned}$$

$$\begin{aligned}\cos 2A &= \cos^2 A - \sin^2 A \\ &= \cos^2 A - (1 - \cos^2 A) \\ &= \cos^2 A - 1 + \cos^2 A \\ &= 2\cos^2 A - 1\end{aligned}$$

To summarise:

$$\sin 2A = 2\sin A \cos A$$

$$\begin{aligned}\cos 2A &= \cos^2 A - \sin^2 A \\ &= 2\cos^2 A - 1 \\ &= 1 - 2\sin^2 A\end{aligned}$$

Example 12: Express the following using double angle formulae:

a) $\sin 2X$

$$= 2 \sin X \cos X$$

c) $\cos 2X$ (sine version)

$$= 1 - 2 \sin^2 X$$

e) $\sin 5Q$

$$= 2 \sin \frac{5}{2} Q \cos \frac{5}{2} Q$$

b) $\sin 6Y$

$$\begin{aligned} &= \sin(3Y + 3Y) \\ &= \sin 3Y \cos 3Y + \cos 3Y \sin 3Y \\ &= 2 \sin 3Y \cos 3Y \end{aligned}$$


d) $\cos 8H$ (cosine version)

$$\begin{aligned} &= \cos(4H + 4H) \\ &= \cos 4H \cos 4H - \sin 4H \sin 4H \\ &= \cos^2 4H - \sin^2 4H \\ &= \cos^2 4H - (1 - \cos^2 4H) \\ &= \cos^2 4H - 1 + \cos^2 4H \\ &= 2 \cos^2 4H - 1 \end{aligned}$$

f) $\cos \theta$ (cos and sin version)

$$= \cos^2 \left(\frac{1}{2} \theta \right) - \sin^2 \left(\frac{1}{2} \theta \right)$$

Example 13: $\sin\theta = \frac{2}{\sqrt{13}}$, where θ is an acute angle. Find the exact values of:



a) $\sin(2\theta)$

$$\begin{aligned} &= 2\sin\theta \cos\theta \\ &= 2 \times \frac{2}{\sqrt{13}} \times \frac{3}{\sqrt{13}} \\ &= \frac{12}{13} \end{aligned}$$

b) $\cos(2\theta)$

$$\begin{aligned} &= \cos^2\theta - \sin^2\theta \\ &= \left(\frac{3}{\sqrt{13}}\right)^2 - \left(\frac{2}{\sqrt{13}}\right)^2 \\ &= \frac{9}{13} - \frac{4}{13} \\ &= \frac{5}{13} \end{aligned}$$

$$\begin{aligned} &2\cos^2\theta - 1 \\ &= 2\left(\frac{3}{\sqrt{13}}\right)^2 - 1 \\ &= 2\left(\frac{9}{13}\right) - 1 \\ &= \frac{18}{13} - \frac{13}{13} \\ &= \frac{5}{13} \end{aligned}$$

$$\begin{aligned} &1 - 2\sin^2\theta \\ &= 1 - 2\left(\frac{2}{\sqrt{13}}\right)^2 \\ &= 1 - 2\left(\frac{4}{13}\right) \\ &= 1 - \frac{8}{13} \\ &= \frac{13}{13} - \frac{8}{13} \\ &= \frac{5}{13} \end{aligned}$$

Ex 11G Q2, 3, 4, 6, 7, 9.

$$s = \frac{D}{t}$$

$$D = st$$

$$\begin{aligned} D &= 112 \times 0.0004 \\ D &= 0.0467 \text{ km} \end{aligned}$$

$$D = \underline{\underline{46.7 \text{ m}}}$$

$$s = 112$$

$$t = 1.5 \text{ s}$$

$$\begin{aligned} 1.5 \div 60 \div 60 \\ &= 0.0004 \text{ h} \end{aligned}$$

Example 14: Prove that $\frac{\sin 2x}{1 + \cos 2x} = \tan x$

$$\begin{aligned}
 & \underline{\text{LHS}} \\
 & \frac{\sin 2x}{1 + \cos 2x} \\
 & = \frac{2 \sin x \cos x}{1 + 2 \cos^2 x - 1} \\
 & = \frac{2 \sin x \cos x}{2 \cos^2 x} \\
 & = \frac{2 \sin x}{2 \cos x} = \frac{\sin x}{\cos x} \\
 & = \underline{\tan x}
 \end{aligned}$$

$$\begin{aligned}
 & \sin 2x \\
 & = 2 \sin x \cos x \\
 & \cos 2x \\
 & = 2 \cos^2 x - 1
 \end{aligned}$$

9.5 Solving non-unitary argument trig equations.

a) Solve for x:

$$2\sin 2x^\circ - 1 = 0 \quad 0 \leq x \leq 360$$

$$2\sin 2x^\circ = 1 \quad 0 \leq x \leq 1080$$

$$\sin 2x^\circ = \frac{1}{2}$$

$$2x^\circ = \sin^{-1}\left(\frac{1}{2}\right)$$

$$2x^\circ = 30^\circ \text{ (R.A.)}$$

$$2x^\circ = 30^\circ, 150^\circ$$

$$x^\circ = 15^\circ, 75^\circ, 195^\circ, 255^\circ$$

180-x	0+x
T	C

b) Solve for x:

$$\sqrt{3} \tan \frac{1}{2}x^\circ = 1 \quad 0 \leq x \leq \pi$$

$$\tan \frac{1}{2}x^\circ = \frac{1}{\sqrt{3}} \quad 0 \leq x \leq 180^\circ$$

$$\frac{1}{2}x^\circ = \tan^{-1}\left(\frac{1}{\sqrt{3}}\right)$$

$$\frac{1}{2}x^\circ = 30^\circ \text{ (R.A.)}$$

$$\frac{1}{2}x^\circ = 30^\circ, 210^\circ$$

$$x^\circ = 60^\circ, 420^\circ$$

~~420~~
out of range.

$$x = 60^\circ$$

$$x = \frac{\pi}{3}$$


180+30	A
T	C

$$\sin x^\circ = -\frac{1}{2}$$

$$x = \sin^{-1}\left(-\frac{1}{2}\right)$$

Ex 4H
Q1 for

Tuesday 11/12/18




Calculus 3: Further Calculus

Let $f(x) = \sin x$ and $g(x) = \cos x$. The graphs of $y = f(x)$ and $y = g(x)$ are shown below, where the x -axis is measured in radians. Tangents to each curve have been drawn at the following points:

On $y = \sin x$, the tangent at $x = 0$ has $m = 1$, and the tangent at $x = \pi$ has $m = -1$.

On $y = \cos x$, the tangent at $x = \frac{\pi}{2}$ has $m = -1$, and the tangent at $x = \frac{3\pi}{2}$ has $m = 1$.

Draw the graphs of $y = f'(x)$ and $y = g'(x)$ below.



If $y = \sin x$, $\frac{dy}{dx} =$

If $y = \cos x$, $\frac{dy}{dx} =$

Example 1: Find the derivative in each case:

a) $y = 4\sin x$

b) $f(x) = 2\cos x$

c) $g(x) = -\frac{1}{2}\cos x$

d) $h = -5\sin x$

As integration is the opposite of differentiation, we can also say that:

$\int \cos x dx =$

$\int \sin x dx =$

Example 2: Find:

a) $\int 24\cos x dx$

$= 24\sin x + C$

b) $\int -3\sin x dx$

$= 3\cos x + C$

c) $\int (3x - \cos x) dx$

$\frac{3x^2}{2} - \sin x + C$

IMPORTANT!

- Definite Integrals of sin and cos functions MUST be done in radians!
- NEVER ignore any brackets where the limit is zero!

Example 3: Evaluate:

$$\begin{aligned}
 \text{a) } \int_0^{\pi/2} \sin x \, dx &= [-\cos x]_0^{\pi/2} \\
 &= [-\cos \frac{\pi}{2}] - [-\cos 0] \\
 &= 0 + 1 \\
 &= \underline{\underline{1}}.
 \end{aligned}$$

$$\begin{aligned}
 \text{c) } \int_0^3 2 \cos x \, dx &= [2 \sin x]_0^3 \\
 &= [2 \times \sin 3] - [2 \times \sin 0] \\
 &= [0.28] - [0] \\
 &= \underline{\underline{0.28}}.
 \end{aligned}$$

$$\begin{aligned}
 \text{b) } \int_0^{\pi/4} (\sin x - \cos x) \, dx &= [-\cos x - \sin x]_0^{\pi/4} \\
 &= [-\cos \frac{\pi}{4} - \sin \frac{\pi}{4}] \\
 &\quad - [-\cos 0 - \sin 0] \\
 &= [-\frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}}] - [-1] \\
 &= \underline{\underline{\frac{1}{\sqrt{2}} + 1}}.
 \end{aligned}$$

The Chain Rule

Example 4: By first expanding the brackets, find the derivatives of the following functions:

a) $y = (3x + 1)^2$

$$\frac{dy}{dx} = 2(3x+1)^1 \times 3$$

$$= 6(3x+1)$$

$$\therefore \frac{dy}{dx} = \underline{\quad}(3x + 1) \times \underline{\quad}$$

b) $y = (2x^2 - 1)^2$

$$\frac{dy}{dx} = 2(2x^2-1)^1 \times 4x$$

$$= 8x(2x^2-1)$$

$$\therefore \frac{dy}{dx} = \underline{\quad}(2x^2 - 1) \times \underline{\quad}$$

c) $y = (2x + 1)^3$

$$\frac{dy}{dx} = 3(2x+1)^2 \times 2$$

$$= \underline{\underline{6(2x+1)^2}}$$

$$\therefore \frac{dy}{dx} = \underline{\quad}(2x + 1)^2 \times \underline{\quad}$$

In each case, we can factorise the answer to give us back the original function, which has been differentiated as if it was just an x^2 or x^3 term (multiply by the old power, drop the power by one), and then multiplied by the derivative of the function in the bracket.

This is known as the **Chain Rule**, and can be written generally for brackets with powers as:

For $f(x) = a (\dots)^n$, $f'(x) = an (\dots)^{n-1} \times (\text{DOB})$
 where DOB = the Derivative Of the Bracket

Example 6: Differentiate:

a) $y = \sin(3x)$

$$\frac{dy}{dx} = \cos 3x \times 3$$

$$= \underline{\underline{3 \cos 3x}}$$

b) $f(x) = \cos\left(\frac{\pi}{4} - 2x\right)$

$$f'(x) = -\sin\left(\frac{\pi}{4} - 2x\right) \times (-2)$$

$$= 2 \sin\left(\frac{\pi}{4} - 2x\right)$$

c) $y = \sin(x^2)$

$$\frac{dy}{dx} = \cos(x^2) \times 2x$$

$$= \underline{\underline{2x \cos x^2}}$$

Example 7: Find the equation of the tangent to $y = \sin\left(2x + \frac{\pi}{3}\right)$ when $x = \frac{\pi}{6}$.

$$\frac{dy}{dx} = \cos\left(2x + \frac{\pi}{3}\right) \times 2$$

$$= 2 \cos\left(2x + \frac{\pi}{3}\right)$$

When $x = \frac{\pi}{6}$

$$\frac{dy}{dx} = 2 \cos\left(2 \times \frac{\pi}{6} + \frac{\pi}{3}\right)$$

$$= 2 \cos \frac{2\pi}{3}$$

$$= 2 \times \left(-\frac{1}{2}\right)$$

$$= \underline{\underline{-1}}$$

$m = -1$

$$y = \sin\left(\frac{2\pi}{3}\right)$$

$$y = \frac{\sqrt{3}}{2}$$

$$\left(\frac{\pi}{6}, \frac{\sqrt{3}}{2}\right)$$

$$y - b = m(x - a)$$

$$y - \frac{\sqrt{3}}{2} = -1 \left(x - \frac{\pi}{6}\right)$$

$$y - \frac{\sqrt{3}}{2} = -x + \frac{\pi}{6}$$

$$y = -x + \frac{\pi}{6} + \frac{\sqrt{3}}{2}$$

Further Integration

We have seen that integration is **anti-differentiation**, i.e. the opposite of differentiating.

As finding the derivative of a function with a bracket included multiplying by DOB, then integrating must also include **dividing** by DOB.

To integrate:

$$\int (ax + b)^n dx = \frac{(ax + b)^{n+1}}{(n+1) \times a} + C$$

Important Point: Integration is more complicated than differentiation!

This method only works for **linear** functions inside the bracket, i.e. the highest power = 1. To find, e.g., $\int (g^3 - 7)^2 dg$, we would have to multiply out the bracket and integrate each term separately.

Example 8: Evaluate:

<p>a) $\int (x+3)^3 dx$</p> $= \frac{(x+3)^4}{4 \times 1} + C.$ $= \frac{1}{4} (x+3)^4 + C$ <hr/>	<p>b) $\int (4x-7)^9 dx$</p> $= \frac{(4x-7)^{10}}{10 \times 4} + C$ $= \frac{1}{40} (4x-7)^{10} + C$ <hr/>	<p>c) $\int \frac{dt}{(4t+9)^2} \left(t \neq -\frac{9}{4} \right)$</p> $= \int \frac{1}{(4t+9)^2} dt$ $= \int (4t+9)^{-2} dt.$ $= \frac{(4t+9)^{-1}}{-1 \times 4} + C.$ $= -\frac{1}{4} (4t+9)^{-1} + C$ <hr/>
--	--	--

$$d) \int_1^2 (2t+5)^3 dt$$

$$= \left[\frac{(2t+5)^4}{4 \times 2} \right]_1^2$$

$$= \left[\frac{(2 \times 2 + 5)^4}{8} \right] - \left[\frac{(2 \times 1 + 5)^4}{8} \right]$$

$$= 820 \cdot 125 - 300 \cdot 125$$

$$= \underline{\underline{520}}$$

$$e) \int_0^6 \frac{dx}{\sqrt{4x+1}} \quad \left(x > -\frac{1}{4} \right)$$

$$= \int_0^6 \frac{1}{\sqrt{4x+1}} dx$$

$$= \int_0^6 \frac{1}{(4x+1)^{1/2}} dx$$

$$= \int_0^6 (4x+1)^{-1/2} dx$$

$$= \left[\frac{(4x+1)^{1/2}}{\frac{1}{2} \times 4} \right]_0^6$$

$$= \left[\frac{\sqrt{4x+1}}{2} \right]_0^6$$

$$= \left[\frac{\sqrt{4 \times 6 + 1}}{2} \right] - \left[\frac{\sqrt{4 \times 0 + 1}}{2} \right]$$

$$= \left[\frac{\sqrt{25}}{2} \right] - \left[\frac{1}{2} \right]$$

For functions including sine and cosine components:

$$\int \sin(ax+b)dx = -\frac{1}{a}\cos(ax+b)+C$$

$$\int \cos(ax+b)dx = \frac{1}{a}\sin(ax+b)+C$$

Example 9: Evaluate:

a) $\int \sin 4x dx$

$$= -\frac{\cos 4x}{4} + C.$$

b) $\int 3\cos 2x dx$

$$= \frac{3\sin 2x}{2} + C$$

c) $\int \sin(1-2x)dx$

$$= -\frac{\cos(1-2x)}{-2} + C$$

$$= \frac{\cos(1-2x)}{2} + C.$$

d) (i) Write $\cos^2 x$ in terms of $\cos 2x$

$$\begin{aligned} \cos 2x &= 2\cos^2 x - 1 \\ 2\cos^2 x - 1 &= \cos 2x \\ 2\cos^2 x &= \cos 2x + 1 \\ \cos^2 x &= \frac{1}{2}(\cos 2x + 1) \\ &= \frac{1}{2}\cos 2x + \frac{1}{2}. \end{aligned}$$


(ii) Hence find $\int 4\cos^2 x \, dx$

$$\begin{aligned} &= \int 4\left(\frac{1}{2}\cos 2x + \frac{1}{2}\right) dx. \\ &= \int 2\cos 2x + 2 \, dx. \\ &= \frac{2\sin 2x}{2} + 2x + C \\ &= \sin 2x + 2x + C \end{aligned}$$

e) Evaluate $\int_0^{2\pi} \sin\left(\frac{1}{2}x\right) dx$

$$\begin{aligned} &= \left[\frac{-\cos\left(\frac{1}{2}x\right)}{\frac{1}{2}} \right]_0^{2\pi} \\ &= \left[-2\cos\frac{1}{2}x \right]_0^{2\pi} \\ &= \left[-2\cos\pi \right] - \left[-2\cos 0 \right] \\ &= \left[-2 \times (-1) \right] - \left[-2 \right] \\ &= 2 + 2 \\ &= \underline{\underline{4}}. \end{aligned}$$

Example 10: Find the area enclosed by $y = \sin\left(2x - \frac{\pi}{4}\right)$, the x-axis and the lines $x = 0$ and $x = \frac{\pi}{2}$.



Calcs x-axis

when $y = 0$

$$\sin\left(2x - \frac{\pi}{4}\right) = 0$$

$$2x - \frac{\pi}{4} = \sin^{-1}(0)$$

$$2x - \frac{\pi}{4} = 0, \pi, 2\pi$$

$$2x = \frac{\pi}{4}, \frac{5\pi}{4}, \frac{9\pi}{4}$$

$$x = \frac{\pi}{8}, \frac{5\pi}{8}, \frac{9\pi}{8}$$

$$\int_0^{\pi/8} \sin\left(2x - \frac{\pi}{4}\right) dx$$

$$= \left[-\frac{\cos\left(2x - \frac{\pi}{4}\right)}{2} \right]_0^{\pi/8}$$

$$= -\frac{\cos(0)}{2} - \left[-\frac{\cos\left(-\frac{\pi}{4}\right)}{2} \right]$$

$$= -\frac{1}{2} - \frac{-\frac{1}{\sqrt{2}}}{2}$$

$$= -\frac{1}{2} + \frac{1}{2\sqrt{2}}$$

$$= \frac{-2 + \sqrt{2}}{4} \times (-)$$

$$\int_{\pi/8}^{\pi/2} \sin\left(2x - \frac{\pi}{4}\right) dx$$

$$= \left[-\frac{\cos\left(2x - \frac{\pi}{4}\right)}{2} \right]_{\pi/8}^{\pi/2}$$

$$= \left[-\frac{\cos\left(\pi - \frac{\pi}{4}\right)}{2} \right] - \left[-\frac{\cos(0)}{2} \right]$$

$$= \frac{\frac{\sqrt{2}}{2}}{2} + \frac{1}{2}$$

$$= \frac{\sqrt{2}}{4} + \frac{1}{2}$$

$$= \frac{2 + \sqrt{2}}{4}$$

$$A_T = \frac{2 - \sqrt{2}}{4} + \frac{2 + \sqrt{2}}{4}$$

$$A_T = \frac{4}{4} = 1 \text{ unit}^2$$

Uses of Calculus in Real Life Situations

In the same way that geometry is the study of shape, calculus is the study of how functions change. This means that wherever a system can be described mathematically using a function, calculus can be used to find the ideal conditions (as we have seen using optimisation) or to use the rate of change at a given time to find the total change (using integration).

As a result, calculus is used throughout the sciences: in Physics (Newton's Laws of Motion, Einstein's Theory of Relativity), Chemistry (reaction rates, radioactive decay), Biology (modelling changes in population), Medicine (using the decay of drugs in the bloodstream to determine dosages), Economics (finding the maximum profit), Engineering (maximising the strength of a building whilst using the minimum of material, working out the curved path of a rocket in space) and more.

Example 11: In Physics, the formulae for kinetic energy (E_k) and momentum (p) are respectively.

$$E_k = \frac{1}{2}mv^2 \quad \text{and} \quad p = mv$$

a) How could the formula for momentum be obtained from the formula for kinetic energy?

$$\begin{aligned} E_k' &= mv \\ &= p. \\ \text{Differentiate} \end{aligned}$$

b) How could the formula for kinetic energy be obtained from the formula for momentum?


$$\begin{aligned} \int mv &= \frac{1}{2}mv^2 + C. \\ \int 3v &= \frac{3v^2}{2} \end{aligned}$$

Displacement, Velocity and Acceleration

The most common use of this approach considers the link between displacement, velocity and acceleration.

When an object moves on a journey, we normally think of the total distance travelled.

Displacement is the straight line distance between the start and end points of a journey (so the displacement is not necessarily the same as the distance travelled!)



As displacement is a “straight-line” measurement, it involves direction and therefore is a **vector** quantity: another name for displacement is the **position**.


Velocity is the vector equivalent of speed, i.e. if speed is a measure of the distance travelled in a given time, then velocity is a measure of the change in displacement which occurs in a given time.

Velocity is defined as the **rate of change of displacement with respect to time**.

Acceleration measures the change in velocity of an object in a given time: if two race cars have the same top speed, then the one which can get to that top speed first would win a race.

Acceleration is defined as the **rate of change of velocity with respect to time**.

If one of either displacement, velocity or acceleration can be described using a function, then the other two can be obtained using either differentiation or integration, i.e.:



Example 12: The displacement s cm at a time t seconds of a particle moving in a straight line is given by the formula $s = t^3 - 2t^2 + 3t$.

a) Find its velocity v cm/s after 3 seconds.

$$\begin{aligned}\frac{ds}{dt} &= 3t^2 - 4t + 3 \\ v &= 3(3)^2 - 4(3) + 3 \\ v &= 27 - 12 + 3 \\ v &= \underline{18 \text{ cm/s}}.\end{aligned}$$

b) The time at which its acceleration a is equal to 26 cm/s^2 .

$$\begin{aligned}v &= 3t^2 - 4t + 3 \\ \frac{dv}{dt} &= 6t - 4 \\ a &= 6t - 4 \\ 26 &= 6t - 4 \\ 6t &= 30 \\ t &= \underline{5}\end{aligned}$$

Example 13: The velocity of an electron is given by the formula $v(t) = 5 \sin\left(2t - \frac{\pi}{4}\right)$.

- a) Find the first time when its acceleration is at its maximum. b) Find a formula for the displacement of the electron, given that $s = 0$ when $t = 0$.

$$v'(t) = 5 \cos\left(2t - \frac{\pi}{4}\right) \times 2$$

$$a = 10 \cos\left(2t - \frac{\pi}{4}\right)$$

SP's when $v'(t) = 0$.

$$10 \cos\left(2t - \frac{\pi}{4}\right) = 0$$

$$\cos\left(2t - \frac{\pi}{4}\right) = 0$$

$$2t - \frac{\pi}{4} = \frac{\pi}{2}, \frac{3\pi}{2}$$

$$2t = \frac{3\pi}{4}, \frac{7\pi}{4}$$

$$t = \frac{3\pi}{8}, \frac{7\pi}{8}$$

t	0	$\frac{3\pi}{8}$	$\frac{7\pi}{8}$	π
$v'(t)$	$+$	0	-0	$+$
Shape	$ $	$-$	$ $	$ $
		Max when $t = \frac{3\pi}{8}$		

$$\int 5 \sin\left(2t - \frac{\pi}{4}\right) dt$$

$$= \frac{-5 \cos\left(2t - \frac{\pi}{4}\right)}{2} + C$$

$$s = \frac{-5}{2} \cos\left(2t - \frac{\pi}{4}\right) + C$$

$$0 = \frac{-5}{2} \times \cos\left(-\frac{\pi}{4}\right) + C$$

$$0 = -2.5 + C$$

$$C = 2.5$$


$$s = \frac{-5}{2} \cos\left(2t - \frac{\pi}{4}\right) + 2.5$$

Past Paper Example 1: A curve has equation $y = (2x - 9)^{\frac{1}{2}}$. Part of the curve is shown in the diagram opposite.

- a) Show that the tangent to the curve at the point where $x = 9$ has equation $y = \frac{1}{3}x$.

$$\begin{aligned} \frac{dy}{dx} &= \frac{1}{2} (2x - 9)^{-\frac{1}{2}} \times 2 \\ &= \frac{1}{\sqrt{2x - 9}} \\ \text{When } x &= 9 \quad \frac{1}{\sqrt{2 \times 9 - 9}} = \frac{1}{3} \end{aligned}$$

- b) Find the coordinates of A, and hence find the shaded area.



$$\begin{aligned} \text{When } x &= 9 \\ y &= (2 \times 9 - 9)^{\frac{1}{2}} \\ y &= 3 \\ &(9, 3) \\ y - b &= m(x - a) \\ y - 3 &= \frac{1}{3}(x - 9) \\ y &= \frac{1}{3}x \end{aligned}$$

Past Paper Example 2: A curve for which $\frac{dy}{dx} = 3\sin 2x$ passes through the point $(\frac{5\pi}{12}, \sqrt{3})$.

Find y in terms of x .

$$y = -\frac{3 \overset{\text{Cos}}{\cancel{\sin}}(2x)}{2} + C.$$

When $x = \frac{5\pi}{12}$ $y = \sqrt{3}$.

$$\sqrt{3} = -\frac{3 \overset{\text{Cos}}{\cancel{\sin}} \frac{5\pi}{12}}{2} + C.$$

$$\sqrt{3} = \frac{3\sqrt{3}}{4} + C$$

$$C = \sqrt{3} - \frac{3\sqrt{3}}{4}$$

$$C = \frac{\sqrt{3}}{4}$$

$$y = -\frac{3 \text{Cos } 2x}{2} + \frac{\sqrt{3}}{4}.$$

Past Paper Example 3: Find the values of x for which the function $f(x) = 2x + 3 + \frac{18}{x-4}$, $x \neq 4$, is increasing.

functions increase when $f'(x) > 0$.

$$f(x) = 2x + 3 + 18(x-4)^{-1}$$

$$f'(x) = 2 - 18(x-4)^{-2} \times 1$$

$$= 2 - \frac{18}{(x-4)^2}$$

$$2 - \frac{18}{(x-4)^2} > 0 \quad \times (x-4)^2$$


$$2(x-4)^2 - 18 > 0$$

$$2(x-4)^2 > 18$$

$$(x-4)^2 > 9$$

$$x < 1, x > 7$$

$$\begin{aligned} (x-4)^2 &= 9 \\ (x-4)^2 - 9 &= 0 \\ x^2 - 8x + 16 - 9 &= 0 \\ x^2 - 8x + 7 &= 0 \end{aligned}$$




$$(x-7)(x-1) = 0$$

$$x = 7, x = 1$$

Trigonometry: The Wave Function

It is possible to model the behaviour of waves in real-life situations (e.g. the interaction of sound waves or the tides where two bodies of water meet) using trigonometry. Consider the result of combining the waves represented by the functions $y = \sin x^\circ$ and $y = \cos x^\circ$. To find what the resultant graph would look like, complete the table of values (accurate to 1 d.p.) and plot on the axes below.

	0°	45°	90°	135°	180°	225°	270°	315°	360°
$\sin x^\circ$	0	0.7	1	0.7	0	-0.7	-1	-0.7	0
$\cos x^\circ$	1	0.7	0	-0.7	-1	-0.7	0	0.7	1
$\sin x^\circ + \cos x^\circ$	1	1.4	1	0	-1	-1.4	-1	0	1



Max = 1.4

Min = -1.4

Max when $x =$ 45°

Min when $x =$ 225°

$\therefore y = 1.4 \cos(x - 45)^\circ$

$y = \sqrt{2} \cos(x - 45^\circ)$

Looking at the graph of $y = \sin x^\circ + \cos x^\circ$ above, we can compare it to cosine graph shifted 45° to the right (i.e. $y = \cos(x - \alpha)^\circ$), and stretched vertically by a factor of roughly 1.4 (i.e. $y = k \cos x^\circ$).

It is important to note, however, that the graph could also be described as a cosine graph shifted to the *left*, and also as a sine graph! Therefore, $y = \sin x^\circ + \cos x^\circ$ could also be written as:

$$y = 1.4 \cos(x + \underline{315}) \quad \text{OR} \quad y = 1.4 \sin(x - \underline{315}) \quad \text{OR} \quad y = 1.4 \sin(x + \underline{45})$$

Rather than drawing an approximate graph, it is more useful if we use an algebraic method.

NOTE: you will only be asked to use one specific form to describe a function, not all four!

Example 1: Write $\sin x^\circ + \cos x^\circ$ in the form $k \cos(x - \alpha)^\circ$, where $0 \leq \alpha \leq 360$.

Handwritten solution for Example 1:

$$R \sin \alpha = 1$$

$$R \cos \alpha = 1$$

$$\frac{R \sin \alpha}{R \cos \alpha} = 1$$

$$\tan \alpha = 1$$

$$\alpha = \tan^{-1}(1)$$

$$\alpha = \underline{\underline{45^\circ}} \text{ (R.A.)}$$

$$\alpha = 45^\circ$$

$$\sqrt{2} \cos(x - 45^\circ)$$

Diagrammatic solution:

$$R \cos(x - \alpha)$$

$$R (\cos x \cos \alpha + \sin x \sin \alpha)$$

$$R \cos x \cos \alpha + R \sin x \sin \alpha$$

$$\textcircled{1} \quad \underline{R \cos x \cos \alpha} + \underline{R \sin x \sin \alpha}$$

Signs diagram:

S ✓	✓ ✓
✓	✓


Labels: T, A, C

Lesson Starter - 5B1 - Mon 4/3/19

1) Differentiate the following function.

$$f(x) = \frac{x^3 - 2x^2 + x - 3}{x^2}$$

2) Find the equation of the line BD, the median of AC



$$m_{BD} = \frac{8-7}{1-3}$$

$$= \frac{1}{-2}$$

$$MP_{AC} = (3, 7)$$

$$y - b = m(x - a)$$

$$y - 7 = -\frac{1}{2}(x - 3)$$

$$2y - 14 = -x + 3$$

$$\underline{2y + x - 17 = 0}$$

This technique can also include the difference between waves and to include double (or higher) angles, but only when the angles of both the sin and cos term are the same (i.e. $2\cos 2x + 5\sin 2x$ can be written as a wave function, but $2\cos 2x + 5\sin 3x$ could not).

Example 2: Write $\sin x - \sqrt{3}\cos x$ in the form $k \cos(x - \alpha)$, where $0 \leq \alpha \leq 2\pi$

$$R \sin x = 1$$

$$k \cos x = -\sqrt{3}$$

$$k = \sqrt{1^2 + (-\sqrt{3})^2}$$

$$k = \sqrt{1 + 3}$$

$$k = 2$$

$$\frac{\sin \alpha}{\cos \alpha} = \frac{1}{-\sqrt{3}}$$

$$\tan \alpha = -\frac{1}{\sqrt{3}}$$

$$\alpha = \tan^{-1}\left(\frac{1}{\sqrt{3}}\right)$$

$$\alpha = \frac{\pi}{6} \text{ (P.A.)}$$

$$\alpha = \pi - \frac{\pi}{6}$$

$$\alpha = \frac{5\pi}{6}$$


$$k \cos(x - \alpha)$$

$$= k(\cos x \cos \alpha + \sin x \sin \alpha)$$

$$= k \cos x \cos \alpha + k \sin x \sin \alpha$$

$$= k \cos x \cos \alpha + k \sin x \sin \alpha$$

①
②



$$\therefore \underline{2 \cos\left(x - \frac{5\pi}{6}\right)}$$

Example 3: Write $12 \cos x - 5 \sin x$ in the form $k \sin(x - \alpha)$, where $0 \leq \alpha \leq 360$


$$\begin{aligned} -k \sin \alpha &= 12 \\ k \sin \alpha &= -12 \\ k \cos \alpha &= -5 \end{aligned}$$

$$\begin{aligned} &k(\sin x \cos \alpha - \cos x \sin \alpha) \\ &k \sin x \cos \alpha - k \cos x \sin \alpha \\ &k \cos \alpha \sin x - k \sin \alpha \cos x \end{aligned}$$

$$\begin{aligned} k &= \sqrt{(-12)^2 + (-5)^2} \\ k &= \sqrt{144 + 25} \\ k &= \sqrt{169} \\ k &= 13 \end{aligned}$$

$$\begin{aligned} \tan \alpha &= \frac{-12}{-5} \\ \tan \alpha &= \frac{12}{5} \\ \alpha &= \tan^{-1}\left(\frac{12}{5}\right) \\ \alpha &= 67.4^\circ \end{aligned}$$

$$\begin{aligned} \alpha &= 180 + 67.4^\circ \\ \alpha &= 247.4^\circ \\ \therefore 13 \sin(x - 247.4^\circ) \end{aligned}$$



Example 4: Write $2\sin 2\theta - \cos 2\theta$ in the form $k \sin(2\theta + \alpha)$, where $0 \leq \alpha < 2\pi$

$$k \sin \alpha = -1$$

$$k \cos \alpha = 2$$

$$k = \sqrt{-1^2 + 2^2}$$

$$k = \sqrt{1 + 4}$$

$$k = \sqrt{5}$$

$$\tan \alpha = -\frac{1}{2}$$

$$\alpha = \tan^{-1}\left(\frac{1}{2}\right)$$

$$\alpha = 26.6^\circ \text{ (D.A.)}$$

$$\frac{360}{-26.6}$$

$$\underline{\underline{333.4}}$$

~~$$\sqrt{5} \sin(2\theta - 26.6^\circ)$$~~

$$\sqrt{5} \sin(2\theta + 333.4^\circ)$$

$$\sqrt{5} \sin(2\theta + 5.82)$$


$$\frac{333.4 \times \pi}{180}$$

$$= 5.82$$

$$k(\sin 2\theta \cos \alpha + \cos 2\theta \sin \alpha)$$

$$= k \cos \alpha \sin 2\theta + k \sin \alpha \cos 2\theta$$

(2) (-1)



Solving Trig Equations Using the Wave Function

In almost all cases, questions like these will be split into two parts, with a) being a “write in the form $y = k \cos(x - \alpha)$ ” followed by b) asking “hence or otherwise solve.....”.

Use the wave function from part a) to solve the equation!

Example 5:

a) Write $2\cos x^\circ - \sin x^\circ$ in the form $k \cos(x - \alpha)^\circ$ where $0 \leq \alpha \leq 360$

b) Hence solve $2\cos x^\circ - \sin x^\circ = -1$ where $0 \leq x \leq 360$

Example 6:

a) Write $\sqrt{3}\sin x + \cos x$ in the form $k \cos(x - \alpha)^\circ$, where $0 \leq \alpha \leq 360^\circ$


$$\begin{aligned}
 k \sin \alpha &= \sqrt{3} \\
 k \cos \alpha &= 1 \\
 k &= \sqrt{3^2 + 1^2} = 2 \\
 \alpha &= 60^\circ \text{ (QA)}
 \end{aligned}$$

$$\begin{aligned}
 k(\cos x \cos \alpha + \sin x \sin \alpha) \\
 k \cos \alpha \cos x + k \sin \alpha \sin x
 \end{aligned}$$



$$\therefore 2 \cos(x - 60)^\circ$$

b) Find algebraically for $0 \leq x \leq 360^\circ$:

(i) The maximum and minimum turning points of $y = \sqrt{3}\sin x + \cos x$.



$$\begin{aligned}
 \text{Max occurs at } 2 \text{ when } x &= 60^\circ \\
 \text{Min occurs at } -2 \text{ when } x &= 240^\circ \\
 \text{Max } (60^\circ, 2) \\
 \text{Min } (240^\circ, -2)
 \end{aligned}$$




(ii) The points of intersection of $y = \sqrt{3}\sin x + \cos x$ with the coordinate axes.

Cuts x-axis @ $(150, 0)$ & $(330, 0)$

Cuts y-axis when $x = 0$.

$$\begin{aligned}
 y &= 2 \cos(0 - 60)^\circ \\
 y &= 2 \cos(-60)^\circ \\
 y &= 2 \times \cos 300^\circ \\
 y &= 2 \times \cos 60^\circ \\
 y &= 2 \times \frac{1}{2} \\
 y &= 1 \\
 \underline{(0, 1)}
 \end{aligned}$$

c) Sketch and annotate the graph of $y = \sqrt{3}\sin x^\circ + \cos x^\circ$ for $0 \leq x \leq 360^\circ$.



Recognising Trig Equations

The trig equations we can be asked to solve at Higher level can be split into three types based on the angle (i.e. x° , $2x^\circ$, $3x^\circ$ etc) and the function(s) (i.e. sin, cos, tan, sin & cos).

<p>Type One: One Function One Angle</p>	<p>e.g.: $2 \sin 4x + 1 = 0$ $\tan^2 x = 3$ $3\sin^2 x - 4\sin x + 1 = 0$</p>	<p>1. Factorise (if necessary) 2. Rearrange to $\sin(\dots) = (\dots)$ [or cos, or tan] 3. Inverse sin/cos/tan to solve</p>
<p>Type Two: Two Functions One Angle</p>	<p>e.g.: $\sin x + \cos x = 1$ $3\cos(2x) + 4 \sin(2x) = 0$ $\cos(4\theta) - \sqrt{3} \sin(4\theta) = -1$</p>	<p>1. Rewrite as a WAVE FUNCTION (choose $k\cos(x - \alpha)$ unless told differently) 2. Solve as Type One</p>
<p>Type Three: Two Angles</p>	<p>e.g.: $5\cos(2\theta) = \cos\theta - 2$ $2\sin(2x) + \sin(x) = -0.5$ $2\cos 2x - \sin x + 5 = 0$</p>	<p>1. Rewrite the double angle and factorise (change $\cos 2x$ to the SINGLE ANGLE function) 2. Solve as Type One</p>

Past Paper Example:

a) The expression $\sqrt{3} \sin x^\circ - \cos x^\circ$ can be written in the form $k \sin(x - \alpha)^\circ$, where $k > 0$ and $0 \leq \alpha < 360$.

Calculate the values of k and α .

$$k(\sin x \cos \alpha - \cos x \sin \alpha)$$

$$k \cos \alpha \sin x - k \sin \alpha \cos x$$

$\sqrt{3}$
 $-$

$-k \sin \alpha = -1$
 $k \sin \alpha = 1$
 $k \cos \alpha = \sqrt{3}$
 $k = \sqrt{(+1)^2 + (\sqrt{3})^2}$
 $k = \sqrt{4}$
 $k = 2$
 $\tan \alpha = \frac{1}{\sqrt{3}}$
 $\alpha = 30^\circ \text{ (1.A.)}$

$2 \sin(x - 30)^\circ$
 $\therefore k = 2$
 $\alpha = 30^\circ$

b) Determine the maximum value of $4 + 5 \cos x^\circ - 5\sqrt{3} \sin x^\circ$, where $0 \leq x < 360$, and state the value of x for which it occurs.

$$\sqrt{3} \sin x - \cos x.$$

$$= 2 \sin(x - 30)^\circ$$


$$5 \cos x - 5\sqrt{3} \sin x.$$

$$5 (\cos x - \sqrt{3} \sin x)$$

$$= -5 (\sqrt{3} \sin x - \cos x).$$

$$= -5 (2 \sin(x - 30)^\circ) + 4.$$

$$= -10 \sin(x - 30)^\circ + 4.$$



$$\text{Max} = 14$$

$$\text{When } x = 300^\circ$$

$$\text{Max @ } 300^\circ$$

$$\text{Max} = 14.$$

Lesson Starter - 5B1 - Mon 18/2/19

1) The point P(5, 2) lies on the curve with equation $y = x^2 - 4x + 7$.

Find the equation of the tangent to the curve at P.

$$\begin{aligned}\frac{dy}{dx} &= 2x - 4 \\ \frac{dy}{dx} &= 2(5) - 4 \\ \frac{dy}{dx} &= 6 \quad m=6\end{aligned}$$

$$\begin{aligned}y - b &= m(x - a) \\ y - 2 &= 6(x - 5) \\ y - 2 &= 6x - 30 \\ y &= 6x - 28\end{aligned}$$

2) Evaluate: $\int_1^2 \frac{1}{6} x^{-2} dx$

$$\begin{aligned}\int \frac{1}{6x^2} &= \left[\frac{\frac{1}{6} x^{-1}}{-1} \right]_1^2 & \int 6x^{-2} \\ &= \left[-\frac{1}{6x} \right]_1^2 & \frac{6x^{-1}}{-1} \\ &= \left[-\frac{1}{12} \right] - \left[-\frac{1}{6} \right] = \frac{1}{12}.\end{aligned}$$

3) a) Solve: $\cos 2x^\circ - 3\cos x^\circ + 2 = 0$ for $0 \leq x < 360^\circ$


$$\begin{aligned}2\cos^2 x - 1 - 3\cos x + 2 &= 0 \\ 2\cos^2 x - 3\cos x + 1 &= 0 \\ \cos x = \frac{1}{2}, \cos x = 1 & \\ x = 60^\circ, 300^\circ, 0^\circ & \\ \text{Let } \cos x = X & \\ 2x^2 - 3x + 1 = 0 & \\ (2x-1)(x-1) = 0 & \\ x = \frac{1}{2}, x = 1 &.\end{aligned}$$

b) Hence solve $\cos 4x^\circ - 3\cos 2x^\circ + 2 = 0$ for $0 \leq x < 360^\circ$

$$\begin{aligned}2x &= 60^\circ, 300^\circ, 240^\circ \\ x &= 30^\circ, 150^\circ, 120^\circ, 180^\circ, 0^\circ, 210^\circ.\end{aligned}$$


Exponential and Logarithmic Functions

Exponential functions are those with variable powers, e.g. $y = a^x$. Their graphs take two forms:



When $a > 1$, the graph:

- is always increasing
- is always positive
- never cuts the x - axis
- passes through (0, 1)
- shows exponential growth



When $0 < a < 1$, the graph

- is always decreasing
- is always positive
- never cuts the x - axis
- passes through (0, 1)
- shows exponential decay

Example 1: Ulanda's population in 2018 was 100 million and it was growing at 6% per annum.

a) Find a formula P_n for the population in millions, n years later.

$$P_n = 100 \times 1.06^n$$

b) Estimate the population in the year 2026

$$P_8 = 100 \times 1.06^8$$

$$= \underline{159 \text{ million}}$$

Example 2: 8000 gallons of oil are lost in an oil spill in Blue Sky Bay. At the beginning of each week a filter plant removes 67% of the oil present.

a) Find a formula G_n for the amount of oil left in the bay after n weeks.

$$G_n = 8000 \times 0.33^n$$

b) After how many complete weeks will there be less than 10 gallons left?

$$G_1 = 8000 \times 0.33^1$$

$$= 2640$$

$$G_2 = 2640 \times 0.33^1$$

$$= 871.2$$

$$G_3 = 871.2 \times 0.33$$

$$= 287.5$$

$$G_4 = 287.5 \times 0.33$$

$$= 94.9$$

Now do Ex 15C Q1 - 4, 7, 8

$$8000 \times 0.33^6 = 10.3$$

$$8000 \times 0.33^7 = 3.4$$

$$G_5 = 94.9 \times 0.33$$

$$= 31.3$$

$$G_6 = 31.3 \times 0.33$$


$$= 10.3$$

$$G_7 = 10.3 \times 0.33$$

$$= 3.4$$

After 7 weeks

Logarithmic Functions



The inverse of an exponential function is known as a **logarithmic** function.

If $f(x) = a^x$, then $f^{-1}(x) = \log_a x$
 (“log to the base a of x ”)

We have seen that the graph of the inverse of a function can be obtained by reflection in the line $y = x$.

Since the graph of $y = 2^x$ passes through the points $(0, 1)$ and $(1, 2)$, then the inverse of $f(x) = 2^x$ must pass through the points $(1, 0)$ and $(2, 1)$.

Example 3: Add the graph of $y = \log_2 x$ to the graph opposite.

Note that:

$y = a^x$ passes through $(0, 1)$ and $(1, a)$
 $y = \log_a x$ passes through $(1, 0)$ and $(a, 1)$

$y = a^x$ means “ a multiplied by itself x times gives y ”

$y = \log_a x$ means “ y is the number of times I multiply a by itself to get x ”

Since the graph does not cross the y -axis, we can only take the logarithm of a positive number

The expression “ $\log_a x$ ” can be read as “ a to the power of what is equal to x ?”, e.g. $\log_2 8$ means “2 to the power of what equals 8?”, so $\log_2 8 = 3$.

Example 4: Write in logarithmic form:

a) $5^2 = 25$
 $\log_5 25 = 2$

b) $12^1 = 12$
 $\log_{12} 12 = 1$

c) $8^{1/3} = 2$
 $\log_8 2 = 1/3$

d) $8^x = y$
 $\log_8 y = x$

e) $1 = q^0$
 $q^0 = 1$
 $\log_q 1 = 0$

f) $(x - 3)^4 = k$
 $\log_{(x-3)} k = 4$

Example 5: Write in exponential form:

a) $3 = \log_5 125$

$$5^3 = 125$$

b) $\log_7 49 = 2$

$$7^2 = 49$$

c) $\log_4 4096 = 6$

$$4^6 = 4096$$

d) $\log_2 \left(\frac{1}{4} \right) = -2$

$$2^{-2} = \frac{1}{4}$$

e) $\log_b g = 5h$

$$b^{5h} = g$$

f) $1 = \log_7 7$

$$7^1 = 7$$

Example 6: Evaluate:

a) $\log_8 64$

$$= 2$$

b) $\log_2 32$

$$= 5$$

c) $\log_{3.5} 3.5$

$$= 1$$

d) $\log_{25} 5$

$$\frac{1}{2}$$

e) $\log_4 \left(\frac{1}{2} \right)$

$$-\frac{1}{2}$$

Since $a^1 = a$, then

$$\log_a a = 1$$

Since $a^0 = 1$, then

$$\log_a 1 = 0.$$

Now do Ex 15E All

Laws of Logarithms

$$\log_a xy = \log_a x + \log_a y$$

$$\log_a \left(\frac{x}{y} \right) = \log_a x - \log_a y$$

$$\log_a x^n = n \log_a x$$

Example 7:

a) $\log_2 4 + \log_2 8 - \log_2 \frac{1}{2}$

b) $2\log_5 10 - \log_5 4$

c) Simplify $\frac{1}{4} (\log_3 810 - \log_3 10)$

Solving Logarithmic Equations

You **MUST** memorise the laws of logarithms to solve log equations! As we can only take logs of **positive** numbers, we must remember to discard any answers which violate this rule!

Example 8: Solve:

$$\text{a) } \log_4 (3x - 2) - \log_4 (x + 1) = \frac{1}{2} \quad \left(x > \frac{2}{3}\right)$$

$$= \log_4 \left(\frac{3x-2}{x+1} \right) = \frac{1}{2}$$

$$\frac{1}{2} = \frac{3x-2}{x+1}$$

$$2 \times \frac{3x-2}{x+1}$$

$$2(x+1) = 3x-2$$

$$2x+2 = 3x-2$$

$$\underline{x = 4}$$

$$\text{b) } \log_6 x + \log_6 (2x - 1) = 2 \quad \left(x > \frac{1}{2}\right)$$

$$\log_6 (x(2x-1)) = 2$$

$$6^2 = x(2x-1)$$

$$2x^2 - x = 36$$

$$2x^2 - x - 36 = 0 \quad \begin{matrix} 2x-9 \\ x+4 \end{matrix}$$

$$(2x-9)(x+4) = 0$$


$$x = \frac{9}{2}, \quad x = -4$$

Discard



The Exponential Function and Natural Logarithms

The graph of the derived function of $y = a^x$ can be plotted and compared with the original function. The new graphs are also exponential functions. Below are the graphs of $y = 2^x$ and $y = 3^x$ (solid lines) and their derived functions (dotted).



The derived function of $y = 2^x$ lies **under** the original graph, but the derived function of $y = 3^x$ lies **above** it.

This means that there must be a value of a between 2 and 3 where the derived function lies **on** the original.

i.e. where $f(x) = f'(x)$

The value of the base of this function is known as e , and is approximately 2.71828.

The function $y = e^x$ is known as The Exponential Function.

The function $y = \log_e x$ is known as the Natural Logarithm of x , and is also written as $\ln x$.

Example 9: Evaluate:

a) e^3
 $= 20.09$

b) $\log_e 120$
 $\log_e = \ln$
 $\ln 120$
 $= 4.79$

Example 10: Solve:

a) $\ln x = 5$
 $\log_e x = 5$
 $x = e^5$
 $x = 148.41$

b) $5^{x-1} = 16$
 $\log_e 5^{(x-1)} = \log_e 16$
 $(x-1) \times \log_e 5 = \log_e 16$
 $x-1 = \frac{\log_e 16}{\log_e 5}$
 $x = \frac{\log_e 16}{\log_e 5} + 1$
 $x = 2.72$

Lesson Starter - 5B1 - Mon 25/2/19

1) Find x when $4\log_x 6 - 2\log_x 4 = 1$

$$\log_x 6^4 - \log_x 4^2 = 1$$

$$\log_x 1296 - \log_x 16 = 1$$

$$\log_x 81 = 1$$

$$x^1 = 81.$$

2) Find: $\int \frac{(x^2 - 2)(x^2 + 2)}{x^2} dx$ 81.

$$= \int \frac{x^4 - 4}{x^2} dx.$$

$$= \int \frac{x^4}{x^2} - \frac{4}{x^2} dx.$$

$$= \int x^2 - 4x^{-2} dx.$$

$$= \frac{x^3}{3} - \frac{4x^{-1}}{-1} + C.$$

$$= \frac{x^3}{3} + 4x^{-1} + C \quad \checkmark \quad \frac{x^3}{3} + \frac{4}{x} + C.$$

Example 11: Atmospheric pressure P_t at various heights above sea level can be determined by using the formula $P_t = P_0 e^{rt}$, where P_0 is the pressure at sea level, t is the height above sea level in thousands of feet, and r is a constant.

a) At 20 000 feet, the air pressure is half that at sea level. Find r accurate to 3 significant figures.

$$\begin{aligned}
 & P_t = P_0 e^{rt} \\
 & P_t = 15 \\
 & P_0 = 30 \\
 & r = ? \\
 & t = 20 \\
 & 15 = 30 \times e^{r \times 20} \\
 & e^{20r} = 0.5 \\
 & \log_e e^{20r} = \log_e 0.5 \\
 & 20r \times \log_e e = \log_e 0.5 \\
 & r = \frac{\log_e 0.5}{20} \\
 & r = -0.0357 \\
 & r \approx -0.0357. \\
 & P_t = P_0 e^{-0.0357t}
 \end{aligned}$$

b) Find the height at which P is 10% of that at sea level.

$$\begin{aligned}
 & P_t = 10 \\
 & P_0 = 100 \\
 & r = -0.0357 \\
 & t = ? \\
 & 10 = 100 \times e^{-0.0357t} \\
 & e^{-0.0357t} = 0.1 \\
 & \log_e e^{-0.0357t} = \log_e 0.1 \\
 & -0.0357t \times \log_e e = \log_e 0.1 \\
 & t = \frac{\log_e 0.1}{-0.0357} \\
 & t = 64.5 \\
 & \text{height} = 64500 \text{ ft}
 \end{aligned}$$

Example 12: A radioactive element decays according to the law $A_t = A_0 e^{kt}$, where A_t is the number of radioactive nuclei present at time t years and A_0 is the initial amount of radioactive nuclei.

a) After 150 years 240g of this material had decayed to 200g.
Find the value of k accurate to 3 s.f.

$$A_t = A_0 e^{kt}$$

$$200 = 240 e^{150k} \quad A_t = 200$$

$$0.83 = e^{150k} \quad A_0 = 240$$

$$(150k) \log_e e = \log_e 0.83 \quad k = ?$$

$$150k = \frac{\log_e 0.83}{\log_e e} \quad t = 150$$

$$k = \underline{\underline{-0.00124}}$$

b) The half-life of the element is the time taken half the mass to decay. Find the half-life of the material.

$$150k = \log_e 0.83$$

$$k = \frac{\log_e 0.83}{150}$$

$$k = \underline{\underline{-0.00124}}$$

$$A_t = A_0 e^{-0.00124t}$$

Example 13: The world population, in billions, t years after 1950 is given by $P = 2.54e^{0.0178t}$.

a) What was the world population in 1950?

When $t = 0$

$$P = 2.54 \times e^0$$

$$P = 2.54$$

$\therefore 2.54$ billion people

b) Find, to the nearest year, the time taken for the population to double.

$$P = 5.08 \text{ billion.}$$

$$5.08 = 2.54 \times e^{0.0178t}$$

$$e^{0.0178t} = 2$$

$$0.0178t = \ln e^2$$


$$t = \frac{\ln e^2}{0.0178}$$

$$t = 38.94 \text{ years.}$$


Using Logs to Analyse Data, Type 1: $y = kx^n \Leftrightarrow \log y = n \log x + \log k$

When the data obtained from an experiment results in an exponential graph of the form $y = kx^n$ as shown below, we can use the laws of logarithms to find the values of k and n .

To begin, take logs of both sides of the exponential equation.




$y = kx^n$



This gives a straight line graph!

Note: the base is not important, as long as the same base is used on both sides.

$y = mx + c$



Example 14: Data are recorded from an experiment and the graph opposite is produced.

a) Find the equation of the line in terms of $\log_{10} x$ and $\log_{10} y$.

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m = \frac{0.903 - 0}{0 - 0.903}$$

$$m = -1$$

$$\log_{10} y = -1 \times \log_{10} x + 0.903$$

$$\log_{10} y = -\log_{10} x + 0.903$$

b) Hence express y in terms of x .

$$\log_{10} y + \log_{10} x = 0.903$$

$$\log_{10}(yx) = 0.903$$


$$yx = 10^{0.903}$$

$$y = \frac{8}{x}$$

Example 15: The data below are plotted and the graph shown is obtained.

x	0	1	2	3	4
$\log_{10} y$	0.602	1.079	1.556	2.033	2.510


a) Express $\log_{10} y$ in terms of x .



b) Hence express y in terms of x .


Related Graphs of Exponentials and Logs

$y = e^x + a$




$y = e^x + a$ is obtained by sliding $y = e^x$:
Vertically upwards if $a > 0$
Vertically downwards if $a < 0$

$y = \ln(x + a)$




$y = \ln(x + a)$ is obtained by sliding $y = \ln x$
Horizontally left if $a > 0$
Horizontally right if $a < 0$

$y = e^{(x+a)}$




$y = e^{(x+a)}$ is obtained by sliding $y = e^x$
Horizontally left if $a > 0$
Horizontally right if $a < 0$

$y = k \ln x$




$y = k \ln x$ is obtained by vertically:
stretching $y = \ln x$ if $k > 1$
compressing $y = \ln x$ if $0 < k < 1$

$y = e^{-x}$




$y = e^{-x}$ is obtained by reflecting $y = e^x$:
in the y -axis

$y = -\ln x$



$y = -\ln x$ is obtained by reflecting $y = \ln x$:
in the x -axis



Example 16: The graph of $y = \log_4 x$ is shown. On the same diagram, sketch:

a) $y = \log_4 4x$

b) $y = \log_4 \left(\frac{1}{4x} \right)$

Past Paper Example 1:


a) Show that $x = 1$ is a root of $x^3 + 8x^2 + 11x - 20 = 0$, and hence factorise $x^3 + 8x^2 + 11x - 20$ fully

b) Solve $\log_2(x + 3) + \log_2(x^2 + 5x - 4) = 3$

Past Paper Example 2: Variables x and y are related by the equation $y = kx^n$.

The graph of $\log_2 y$ against $\log_2 x$ is a straight line through the points $(0, 5)$ and $(4, 7)$, as shown in the diagram.

Find the values of k and n .



Past Paper Example 3: The concentration of the pesticide *Xpesto* in soil is modelled by the equation:

$$P_t = P_0 e^{-kt}$$

P_0 is the initial concentration

where: P_t is the concentration at time t

t is the time, in days, after the application of the pesticide.

a) Once in the soil, the half-life of a pesticide is the time taken for its concentration to be reduced to one half of its initial value.

If the half-life of *Xpesto* is 25 days, find the value of k to 2 significant figures.

b) Eighty days after the initial application, what is the percentage decrease in *Xpesto*?


Past Paper Example 4: Simplify the expression $3\log_e 2e - 2\log_e 3e$ giving your answer in the form $A + \log_e B - \log_e C$, where A, B and C are whole numbers.

$$\begin{aligned}
 & 3\log_e 2e - 2\log_e 3e \\
 & \log_e (2e)^3 - \log_e (3e)^2 \\
 & \log_e 8e^3 - \log_e 9e^2 \\
 & \log_e 8 + \log_e e^3 - (\log_e 9 + \log_e e^2) \\
 & \log_e 8 + \log_e e^3 - \log_e 9 - \log_e e^2 \\
 & \log_e 8 + 3 - \log_e 9 - 2 \\
 & 1 + \log_e 8 - \log_e 9
 \end{aligned}$$

$$\begin{aligned}
 4^x &= 10 & P \times x \\
 \log_e 4^x &= \log_e 10 & P \times x \\
 x \times \log_e 4 &= P \times x \times \log_e 10 \\
 P &= \frac{x \times \log_e 4}{x \times \log_e 10} \\
 P &= 0.6
 \end{aligned}$$

Past Paper Example 5: Two variables x and y satisfy the equation $y = 3(4^x)$.

A graph is drawn of $\log_{10} y$ against x. Show that its equation will be of the form $\log_{10} y = Px + Q$, and state the gradient and y-intercept of this line.



$$\begin{aligned}
 y &= 10^{Px+Q} \\
 y &= 10^{Px} \times 10^Q \\
 y &= (10^4)^x \times 10^3 \\
 10^Q &= 3 \\
 \log_e 10^Q &= \log_e 3 \\
 Q \times \log_e 10 &= \log_e 3 \\
 Q &= \frac{\log_e 3}{\log_e 10} \\
 Q &= 0.48 \\
 \text{gradient} &= 0.6 \\
 \text{y-int} &= 0.48
 \end{aligned}$$